

Review Article

Remarks of Equivalence among Picard, Mann, and Ishikawa Iterations in Normed Spaces

Xue Zhiquan

Received 1 April 2007; Revised 16 April 2007; Accepted 21 June 2007

Recommended by J. R. L. Webb

We show that the convergence of Picard iteration is equivalent to the convergence of Mann iteration schemes for various Zamfirescu operators. Our result extends of Soltuz (2005).

Copyright © 2007 Xue Zhiquan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a real normed space, D a nonempty convex subset of E , and T a self-map of D , let $p_0, u_0, x_0 \in D$. The Picard iteration is defined by

$$p_{n+1} = Tp_n, \quad n \geq 0. \quad (1.1)$$

The Mann iteration is defined by

$$u_{n+1} = (1 - a_n)u_n + a_nTu_n, \quad n \geq 0. \quad (1.2)$$

The Ishikawa iteration is defined by

$$\begin{aligned} y_n &= (1 - b_n)x_n + b_nTx_n, \quad n \geq 0, \\ x_{n+1} &= (1 - a_n)x_n + a_nTy_n, \quad n \geq 0, \end{aligned} \quad (1.3)$$

where $\{a_n\}, \{b_n\}$ are sequences of positive numbers in $[0, 1]$. Obviously, for $a_n = 1$, the Mann iteration (1.2) reduces to the Picard iteration, and for $b_n = 0$, the Ishikawa iteration (1.3) reduces to the Mann iteration (1.2).

Definition 1.1 [1, Definition 1]. Let $T : D \rightarrow D$ be a map for which there exist real numbers a, b, c satisfying $0 < a < 1$, $0 < b < 1/2$, $0 < c < 1/2$. Then T is called a Zamfirescu operator

2 Fixed Point Theory and Applications

if, for each pair x, y in D , T satisfies at least one of the following conditions given in (1)–(3):

- (1) $\|Tx - Ty\| \leq a\|x - y\|$;
- (2) $\|Tx - Ty\| \leq b(\|x - Tx\| + \|y - Ty\|)$;
- (3) $\|Tx - Ty\| \leq c(\|x - Ty\| + \|y - Tx\|)$.

It is easy to show that every Zamfirescu operator T satisfies the inequality

$$\|Tx - Ty\| \leq \delta\|x - y\| + 2\delta\|x - Tx\| \quad (1.4)$$

for all $x, y \in D$, where $\delta = \max\{a, b/(1 - b), c/(1 - c)\}$ with $0 < \delta < 1$ (See Şoltuz [1]). Recently, Şoltuz [1] had studied that the equivalence of convergence for Picard, Mann, and Ishikawa iterations, and proved the following results.

THEOREM 1.2 [1, Theorem 1]. *Let X be a normed space, D a nonempty, convex, closed subset of X , and $T : D \rightarrow D$ an operator satisfying condition Z (Zamfirescu operator). If $u_0 = x_0 \in D$, let $\{u_n\}_{n=0}^{\infty}$ be defined by (1.2) for $u_0 \in D$, and let $\{x_n\}_{n=0}^{\infty}$ be defined by (1.3) for $x_0 \in D$ with $\{a_n\}$ in $[0, 1]$ satisfying $\sum_{n=0}^{\infty} a_n = \infty$. Then the following are equivalent:*

- (i) *the Mann iteration (1.2) converges to the fixed point of T ;*
- (ii) *the Ishikawa iteration (1.3) converges to the fixed point of T .*

THEOREM 1.3 [1, Theorem 2]. *Let X be a normed space, D a nonempty, convex, closed subset of X , and $T : D \rightarrow D$ an operator satisfying condition Z (Zamfirescu operator). If $u_0 = p_0 \in D$, let $\{p_n\}_{n=0}^{\infty}$ be defined by (1.1) for $p_0 \in D$, and let $\{u_n\}_{n=0}^{\infty}$ be defined by (1.2) for $u_0 \in D$ with $\{a_n\}$ in $[0, 1]$ satisfying $\sum_{n=0}^{\infty} a_n = \infty$ and $a_n \rightarrow 0$ as $n \rightarrow \infty$. Then*

- (i) *if the Mann iteration (1.2) converges to x^* and $\lim_{n \rightarrow \infty} (\|u_{n+1} - u_n\|/a_n) = 0$, then the Picard iteration (1.1) converges to x^* ,*
- (ii) *if the Picard iteration (1.1) converges to x^* and $\lim_{n \rightarrow \infty} (\|p_{n+1} - p_n\|/a_n) = 0$, then the Mann iteration (1.2) converges to x^* .*

However, in the above-mentioned theorem, it is unnecessary that, for two conditions, $\lim_{n \rightarrow \infty} (\|u_{n+1} - u_n\|/a_n) = 0$ and $\lim_{n \rightarrow \infty} (\|p_{n+1} - p_n\|/a_n) = 0$. The aim of this paper is to show that the convergence of Picard iteration schemes is equivalent to the convergence of the Mann iteration for Zamfirescu operators in normed spaces. The result improves ones announced by Şoltuz [1, Theorem 2]. We will use a special case of the following lemma.

LEMMA 1.4 [2]. *Let $\{a_n\}$ and $\{\sigma_n\}$ be nonnegative real sequences satisfying the following inequality:*

$$a_{n+1} \leq (1 - \lambda_n)a_n + \sigma_n, \quad (1.5)$$

where $\lambda_n \in (0, 1)$, for all $n \geq n_0$, $\sum_{n=1}^{\infty} \lambda_n = \infty$, and $\sigma_n/\lambda_n \rightarrow 0$ as $n \rightarrow \infty$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

This lemma is apparently due to Vasilen, it is given as [2, Lemma 2.3.6, page 96]. It was rediscovered with a different proof by Weng [3].

2. Main results

THEOREM 2.1. *Let E be a normed space, D a nonempty closed convex subset of E , and $T : D \rightarrow D$ a Zamfirescu operator. Suppose that T has a fixed point $q \in D$. Let $\{p_n\}_{n=0}^\infty$ be defined by (1.1) for $p_0 \in D$, and let $\{u_n\}_{n=0}^\infty$ be defined by (1.2) for $u_0 \in D$ with $\{a_n\}$ in $[0, 1]$ and satisfying $\sum_{n=0}^\infty a_n = \infty$. Then the following are equivalent:*

- (i) *the Picard iteration (1.1) converges to the fixed point of T ;*
- (ii) *the Mann iteration (1.2) converges to the fixed point of T .*

Proof. Let q be a fixed point of T . We will prove (ii) \Rightarrow (i). Suppose that $\|u_n - q\| \rightarrow 0$ as $n \rightarrow \infty$. Applying (1.1) and (1.2), we have

$$\begin{aligned} \|u_{n+1} - p_{n+1}\| &\leq (1 - a_n)\|u_n - Tp_n\| + a_n\|Tu_n - Tp_n\| \\ &\leq (1 - a_n)\|u_n - Tu_n\| + \|Tu_n - Tp_n\|. \end{aligned} \quad (2.1)$$

Using (1.4) with $x = u_n$, $y = p_n$, we have

$$\|Tu_n - Tp_n\| \leq \delta\|u_n - p_n\| + 2\delta\|u_n - Tu_n\|. \quad (2.2)$$

Therefore, from (2.1), we get

$$\begin{aligned} \|u_{n+1} - p_{n+1}\| &\leq \delta\|u_n - p_n\| + (1 - a_n + 2\delta)\|u_n - Tu_n\| \\ &\leq \delta\|u_n - p_n\| + (1 - a_n + 2\delta)(\|u_n - q\| + \|Tu_n - Tq\|) \\ &\leq \delta\|u_n - p_n\| + (1 - a_n + 2\delta)(1 + \delta)\|u_n - q\|, \end{aligned} \quad (2.3)$$

denoted by $A_n = \|u_n - p_n\|$, $\delta = 1 - \lambda$, and $B_n = (1 - a_n + 2\delta)(1 + \delta)\|u_n - q\|$. By Lemma 1.4, we obtain $A_n = \|u_n - p_n\| \rightarrow 0$ as $n \rightarrow \infty$. Hence by $\|p_n - q\| \leq \|u_n - p_n\| + \|u_n - q\|$, we get $\|p_n - q\| \rightarrow 0$ as $n \rightarrow \infty$.

Next, we will prove (i) \Rightarrow (ii), that is, if the Picard iteration converges, then the Mann iteration does too. Now by using Picard iteration (1.1) and Mann iteration (1.2), we have

$$\begin{aligned} \|u_{n+1} - p_{n+1}\| &\leq (1 - a_n)\|u_n - Tp_n\| + a_n\|Tu_n - Tp_n\| \\ &\leq (1 - a_n)\|u_n - p_n\| + (1 - a_n)\|p_n - Tp_n\| + a_n\|Tu_n - Tp_n\| \\ &\leq (1 - a_n)\|u_n - p_n\| + (1 - a_n)(\|p_n - q\| + \|Tp_n - Tq\|) + a_n\|Tu_n - Tp_n\|. \end{aligned} \quad (2.4)$$

On using (1.4) with $x = p_n$, $y = u_n$, we get

$$\begin{aligned} \|Tu_n - Tp_n\| &\leq a_n\delta\|u_n - p_n\| + 2a_n\delta\|p_n - Tp_n\| \\ &\leq a_n\delta\|u_n - p_n\| + 2a_n\delta(\|p_n - q\| + \|Tp_n - Tq\|). \end{aligned} \quad (2.5)$$

4 Fixed Point Theory and Applications

Again, using (1.4) with $x = q$, $y = p_n$, we get

$$\|Tp_n - Tq\| \leq \delta \|p_n - q\|. \quad (2.6)$$

Hence by (2.4)–(2.6), we obtain

$$\begin{aligned} \|u_{n+1} - p_{n+1}\| &\leq (1 - (1 - \delta)a_n) \|u_n - p_n\| + (1 - a_n + 2a_n\delta)(1 + \delta) \|p_n - q\| \\ &\leq (1 - (1 - \delta)a_n) \|u_n - q\| + (1 - a_n + 2a_n\delta)(2 + \delta) \|p_n - q\| \\ &\leq (1 - \lambda a_n)(1 - \lambda a_{n-1}) \|u_{n-1} - q\| + (1 - a_n + 2a_n\delta)(2 + \delta) \|p_n - q\| \\ &\leq (1 - \lambda a_n)(1 - \lambda a_{n-1}) \cdots (1 - \lambda a_0) \|u_0 - q\| \\ &\quad + (1 - a_n + 2a_n\delta)(2 + \delta) \|p_n - q\| \\ &\leq \exp\left(-\lambda \sum_{i=0}^n a_i\right) \|u_0 - q\| + (1 - a_n + 2a_n\delta)(2 + \delta) \|p_n - q\|, \end{aligned} \quad (2.7)$$

where $1 - \delta = \lambda$. Since $\sum_{n=0}^{\infty} a_n = \infty$ and $\|p_n - q\| \rightarrow 0$ as $n \rightarrow \infty$, hence $\|u_n - p_n\| \rightarrow 0$ as $n \rightarrow \infty$. And thus, $\|u_n - q\| \leq \|u_n - p_n\| + \|p_n - q\| \rightarrow 0$ as $n \rightarrow \infty$. This completes the proof. \square

Remark 2.2. Theorem 2.1 improves [1, Theorem 2] in the following sense.

- (1) Both hypotheses $\lim_{n \rightarrow \infty} (\|u_{n+1} - u_n\|/a_n) = 0$ and $\lim_{n \rightarrow \infty} (\|p_{n+1} - p_n\|/a_n) = 0$ have been removed, and the conclusion remains valid.
- (2) The assumption that $u_0 = p_0$ in [1] is superfluous.

THEOREM 2.3. *Let E be a normed space, D a nonempty closed convex subset of E , and $T : D \rightarrow D$ a Zamfirescu operator. Suppose that T has a fixed point $q \in D$. Let $\{p_n\}_{n=0}^{\infty}$ be defined by (1.1) for $p_0 \in D$, and let $\{x_n\}_{n=0}^{\infty}$ be defined by (1.3) for $x_0 \in D$ with $\{a_n\}$ and $\{b_n\}$ in $[0, 1]$ and satisfying $\sum_{n=0}^{\infty} a_n = \infty$. Then the following are equivalent:*

- (i) *the Picard iteration (1.1) converges to the fixed point of T ;*
- (ii) *the Ishikawa iteration (1.3) converges to the fixed point of T .*

Remark 2.4. As previously suggested, Theorem 2.3 reproduces exactly [1, Theorem 1]. Therefore we have the following conclusion: Picard iteration converges to the fixed point of $T \Leftrightarrow$ Mann iteration converges to the fixed point of $T \Leftrightarrow$ Ishikawa iteration converges to the fixed point of T .

Acknowledgments

The author would like to thank the referee and the editor for their careful reading of the manuscript and their many valuable comments and suggestions. The project is supported by the National Science Foundation of China and Shijiazhuang Railway Institute Sciences Foundation.

References

- [1] Ş. M. Şoltuz, “The equivalence of Picard, Mann and Ishikawa iterations dealing with quasi-contractive operators,” *Mathematical Communications*, vol. 10, no. 1, pp. 81–88, 2005.

- [2] F. P. Vasilev, *Numerical Methods for Solving Extremal Problems*, Nauka, Moscow, Russia, 2nd edition, 1988.
- [3] X. Weng, "Fixed point iteration for local strictly pseudo-contractive mapping," *Proceedings of the American Mathematical Society*, vol. 113, no. 3, pp. 727–731, 1991.

Xue Zhiqun: Department of Mathematics and Physics, Shijiazhuang Railway Institute,
Shijiazhuang 050043, China
Email address: xuezhiqun@126.com