

Research Article

# Iterative Algorithms for Nonexpansive Mappings

Yeong-Cheng Liou<sup>1</sup> and Yonghong Yao<sup>2</sup>

<sup>1</sup> Department of Information Management, Cheng Shiu University,  
Kaohsiung 833, Taiwan

<sup>2</sup> Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

Correspondence should be addressed to Yeong-Cheng Liou, simplex\_liou@hotmail.com

Received 13 September 2007; Accepted 25 November 2007

Recommended by Massimo Furi

We suggest and analyze two new iterative algorithms for a nonexpansive mapping  $T$  in Banach spaces. We prove that the proposed iterative algorithms converge strongly to some fixed point of  $T$ .

Copyright © 2008 Y.-C. Liou and Y. Yao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $E$  be a real Banach space,  $C$  a nonempty closed convex subset of  $E$ , and  $T : C \rightarrow C$  a nonexpansive mapping; namely,

$$\|Tx - Ty\| \leq \|x - y\|, \quad (1.1)$$

for all  $x, y \in C$ . We use  $F(T)$  to denote the set of fixed points of  $T$ , that is,  $F(T) = \{x \in C : Tx = x\}$ . Throughout the paper, we assume that  $F(T) \neq \emptyset$ .

Construction of fixed points of nonlinear mappings is an important and active research area. In particular, iterative methods for finding fixed points of nonexpansive mappings have received a vast investigation; see [1–28].

It is well known that the Picard iteration  $x_{n+1} = Tx_n = \dots = T^{n+1}x$  of the mapping  $T$  at a point  $x \in C$  may, in general, not behave well. This means that it may not converge even in the weak topology. One way to overcome this difficulty is to use Mann's iteration method that produces a sequence  $\{x_n\}$  via the recursive manner:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n)Tx_n, \quad n \geq 0, \quad (1.2)$$

where the initial guess  $x_0 \in C$  is chosen arbitrarily. For example, Reich [9] proved that if  $E$  is a uniformly convex Banach space with a Frechet differentiable norm and if  $\{\alpha_n\}$  is chosen such

that  $\sum_{n=1}^{\infty} \alpha_n(1 - \alpha_n) = \infty$ , then the sequence  $\{x_n\}$  defined by (1.2) converges weakly to a fixed point of  $T$ . However, this scheme has only a weak convergence even in a Hilbert space.

Some attempts to construct iteration method so that strong convergence is guaranteed have recently been made. For a sequence  $\{\alpha_n\}$  of real numbers in  $[0, 1]$  and an arbitrary  $u \in C$ , let the sequence  $\{x_n\}$  in  $C$  be iteratively defined by  $x_0 \in C$ ,

$$x_{n+1} = \alpha_n u + (1 - \alpha_n) T x_n. \quad (1.3)$$

The iterative method (1.3) is now referred to as the Halpern iterative method. The interest and importance of Halpern iterative method lie in the fact that strong convergence of the sequence  $\{x_n\}$  is achieved under certain mild conditions on parameter  $\{\alpha_n\}$  in a general Banach space.

Recently, Su and Li [21] introduced the following two new iterative algorithms for a nonexpansive mapping  $T$ : for fixed  $u \in C$ , let the sequences  $\{x_n\}$  and  $\{y_n\}$  be generated by

$$x_{n+1} = \alpha_n u + (1 - \alpha_n) T(\beta_n u + (1 - \beta_n) x_n), \quad (1.4)$$

$$y_{n+1} = \alpha_n(\beta_n u + (1 - \beta_n) T y_n) + (1 - \alpha_n) \frac{1}{n+1} \sum_{i=0}^n T^i y_n, \quad (1.5)$$

respectively. Su and Li proved that the sequences  $\{x_n\}$  and  $\{y_n\}$  converge strongly to some fixed point of  $T$  under some assumptions. Subsequently, Liu and Li [22] extended and improved the corresponding results of Su and Li [21]. But we note that all of the above results have imposed some additional assumptions on parameters. Please see [21, 22] for more details.

Motivated and inspired by the above works, in this paper we construct two new iterative algorithms for approximating fixed points of a nonexpansive mapping  $T$ . We prove that the proposed iterative algorithms converge strongly to a fixed point of  $T$  under some mild conditions.

## 2. Preliminaries

Let  $E$  be a real Banach space with its dual  $E^*$ . Let  $S = \{x \in E : \|x\| = 1\}$  denote the unit sphere of  $E$ . The norm on  $E$  is said to be Gâteaux differentiable if the limit

$$\lim_{t \rightarrow 0} \frac{\|x + t y\| - \|x\|}{t} \quad (2.1)$$

exists for each  $x, y \in S$  and in this case  $E$  is said to be smooth.  $E$  is said to have a uniformly Fréchet differentiable norm if the limit (2.1) is attained uniformly for  $x, y \in S$  and in this case  $E$  is said to be uniformly smooth. It is well known that if  $E$  is uniformly smooth, then the duality map is norm-to-norm uniformly continuous on bounded subsets of  $E$ .

Let  $C \subset E$  be closed convex and  $P$  a mapping of  $E$  onto  $C$ . Then,  $P$  is said to be sunny if  $P(Px + t(x - Px)) = Px$ , for all  $x \in E$  and  $t \geq 0$ . A mapping  $P$  of  $E$  into  $E$  is said to be a retraction if  $P^2 = P$ . If a mapping  $P$  is a retraction, then  $Pz = z$  for every  $z \in R(P)$ , where  $R(P)$  is the range of  $P$ . A subset  $C$  of  $E$  is said to be a sunny nonexpansive retract of  $E$  if there exists a sunny nonexpansive retraction of  $E$  onto  $C$  and it is said to be a nonexpansive retract of  $E$  if there exists a nonexpansive retraction of  $E$  onto  $C$ . If  $E = H$ , the metric projection  $P_C$  is a sunny nonexpansive retraction from  $H$  to any closed convex subset of  $H$ .

We need the following lemmas for proving our main results.

**Lemma 2.1** (see [16]). *Let  $E$  be a real Banach space and  $J$  the normalized duality map on  $E$ . Then, for any given  $x, y \in E$ , the following inequality holds:*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \quad \forall j(x + y) \in J(x + y). \quad (2.2)$$

**Lemma 2.2** (see [20]). *Let  $\{x_n\}$  and  $\{z_n\}$  be bounded sequences in a Banach space  $E$  and let  $\{\alpha_n\}$  be a sequence in  $[0, 1]$  which satisfies the condition  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ . Suppose  $x_{n+1} = \alpha_n x_n + (1 - \alpha_n) z_n$ ,  $n \geq 0$ , and  $\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0$ . Then,  $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$ .*

**Lemma 2.3** (see [17]). *Assume  $\{a_n\}$  is a sequence of nonnegative real numbers such that  $a_{n+1} \leq (1 - b_n)a_n + c_n$ ,  $n \geq 0$ , where  $\{b_n\}$  is a sequence in  $(0, 1)$  and  $\{c_n\}$  is a sequence in  $R$  such that*

- (i)  $\sum_{n=0}^{\infty} b_n = \infty$ ,
- (ii)  $\limsup_{n \rightarrow \infty} c_n/b_n \leq 0$  or  $\sum_{n=0}^{\infty} |c_n| < \infty$ .

*Then,  $\lim_{n \rightarrow \infty} a_n = 0$ .*

**Lemma 2.4** (see [17]). *Let  $C$  be a nonempty closed convex subset of a uniformly smooth Banach space  $E$ . Let  $T : C \rightarrow C$  be a nonexpansive mapping with  $F(T) \neq \emptyset$ . Then, there exists a continuous path  $t \rightarrow z_t$ ,  $0 < t < 1$ , satisfying  $z_t = tu + (1 - t)Tz_t$ , for arbitrary but fixed  $u \in C$ , which converges strongly to a fixed point of  $T$ . Further, if  $Pu = \lim_{t \rightarrow 0} z_t$ , for each  $u \in C$ , then  $P$  is a sunny nonexpansive retraction of  $C$  onto  $F(T)$ .*

**Lemma 2.5** (see [17]). *Let  $C$  be a nonempty closed convex subset of a uniformly smooth Banach space  $E$ . Let  $T : C \rightarrow C$  be a nonexpansive mapping with  $F(T) \neq \emptyset$ . Let  $\{x_n\} \subset C$  be a sequence. Suppose that  $\{x_n\}$  is bounded and  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ . Then,  $\langle u - Pu, j(x_n - Pu) \rangle \leq 0$ .*

### 3. Main results

Let  $C$  be a nonempty closed convex subset of a real uniformly smooth Banach space  $E$ . First, for fixed two different anchors  $u, v \in C$  we define  $x_t = tu + (1 - t)Tx_t$  and  $y_t = tv + (1 - t)Ty_t$ . It follows from Lemma 2.4 that  $x_t$  and  $y_t$  converge strongly to  $Pu$  and  $Pv$ , respectively. Now we assume that anchors  $u$  and  $v$  satisfy condition (P):  $Pu = Pv$ , where  $P$  is a sunny nonexpansive retraction from  $C$  onto  $F(T)$ .

Now we introduce the following iterative algorithm: for fixed  $u, v \in C$  and given  $x_0 \in C$  arbitrarily, find the approximate solution  $\{x_n\}$  by

$$x_{n+1} = \alpha_n u + \beta_n x_n + \gamma_n T(\delta_n v + (1 - \delta_n)x_n), \quad (3.1)$$

where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ , and  $\{\delta_n\}$  are real sequences in  $[0, 1]$ .

*Remark 3.1.* Our iterative algorithm (3.1) can be reviewed as an extension of the iterative algorithm (1.4).

Now we state and prove the strong convergence of the iterative algorithm (3.1).

**Theorem 3.2.** Let  $C$  be a nonempty closed convex subset of a real uniformly smooth Banach space  $E$ . Let  $T : C \rightarrow C$  be a nonexpansive mapping with  $F(T) \neq \emptyset$ . Suppose the sequences  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ , and  $\{\delta_n\}$  in  $(0, 1)$  satisfy  $\alpha_n + \beta_n + \gamma_n = 1$ ,  $n \geq 0$ . Suppose the following conditions are satisfied:

- (i)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (ii)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $\lim_{n \rightarrow \infty} \delta_n = 0$  and  $\delta_n / \alpha_n \leq M$  for some constant  $M \geq 0$ .

For fixed anchors  $u$  and  $v$  satisfying condition (P) and arbitrary given  $x_0 \in C$ , the sequence  $\{x_n\}$  defined by (3.1) converges strongly to  $Pu \in F(T)$ .

*Proof.* Setting  $y_n = \delta_n v + (1 - \delta_n)x_n$  and  $\sigma_n = \alpha_n + \gamma_n \delta_n$ . For  $p \in F(T)$ , we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq \alpha_n \|u - p\| + \beta_n \|x_n - p\| + \gamma_n [\delta_n \|v - p\| + (1 - \delta_n) \|x_n - p\|] \\ &= \alpha_n \|u - p\| + \gamma_n \delta_n \|v - p\| + [\beta_n + \gamma_n (1 - \delta_n)] \|x_n - p\| \\ &\leq \sigma_n \max\{\|v - p\|, \|u - p\|\} + (1 - \sigma_n) \|x_n - p\| \\ &\leq \max\{\|u - p\|, \|v - p\|, \|x_0 - p\|\}, \end{aligned} \tag{3.2}$$

which implies that  $\{x_n\}$  is bounded.

From (3.1), we write  $x_{n+1} = \beta_n x_n + (1 - \beta_n)z_n$ . It is easily seen that

$$\begin{aligned} z_{n+1} - z_n &= \frac{\alpha_{n+1}u + \gamma_{n+1}Ty_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n u + \gamma_n Ty_n}{1 - \beta_n} \\ &= \left( \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) u + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} (Ty_{n+1} - Ty_n) + \left( \frac{\gamma_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_n}{1 - \beta_n} \right) Ty_n. \end{aligned} \tag{3.3}$$

It follows that

$$\begin{aligned} \|z_{n+1} - z_n\| &\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| (\|u\| + \|Ty_n\|) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \|Ty_{n+1} - Ty_n\| \\ &\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| (\|u\| + \|Ty_n\|) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \|y_{n+1} - y_n\|. \end{aligned} \tag{3.4}$$

At the same time, we note that

$$\|y_{n+1} - y_n\| \leq |\delta_{n+1} - \delta_n| (\|v\| + \|x_n\|) + (1 - \delta_{n+1}) \|x_{n+1} - x_n\|. \tag{3.5}$$

It follows from (3.4) and (3.5) that

$$\begin{aligned} \|z_{n+1} - z_n\| &\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| (\|u\| + \|Ty_n\|) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \\ &\quad \times |\delta_{n+1} - \delta_n| (\|v\| + \|x_n\|) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \|x_{n+1} - x_n\|. \end{aligned} \tag{3.6}$$

Then, we have

$$\begin{aligned} \|z_{n+1} - z_n\| &\leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} |(\|u\| + \|Ty_n\|) + \frac{\gamma_{n+1}}{1 - \beta_{n+1}} |\delta_{n+1} - \delta_n| \\ &\quad \times (\|v\| + \|x_n\|) + \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \|x_{n+1} - x_n\| + \|x_{n+1} - x_n\|, \end{aligned} \quad (3.7)$$

which implies that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (3.8)$$

From Lemma 2.2 and (3.8), we obtain  $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$ . Consequently,  $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \beta_n) \|z_n - x_n\| = 0$ .

On the other hand, we observe that

$$\begin{aligned} \|x_n - Tx_n\| &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \alpha_n \|u - Tx_n\| + \beta_n \|x_n - Tx_n\| + \gamma_n \|Ty_n - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \alpha_n \|u - Tx_n\| + \beta_n \|x_n - Tx_n\| + \gamma_n \delta_n \|v - x_n\|, \end{aligned} \quad (3.9)$$

that is,

$$\|x_n - Tx_n\| \leq \frac{1}{1 - \beta_n} \{ \|x_{n+1} - x_n\| + \alpha_n \|u - Tx_n\| + \gamma_n \delta_n \|v - x_n\| \} \rightarrow 0. \quad (3.10)$$

By Lemma 2.5 and (3.10), we can obtain

$$\limsup_{n \rightarrow \infty} \langle u - Pu, j(x_n - Pu) \rangle \leq 0, \quad \limsup_{n \rightarrow \infty} \langle v - Pv, j(x_n - Pv) \rangle \leq 0. \quad (3.11)$$

Since  $\|y_n - x_n\| = \delta_n \|v - x_n\| \rightarrow 0$  and  $Pu = Pv$ , we have

$$\limsup_{n \rightarrow \infty} \langle u - Pu, j(x_n - Pu) \rangle \leq 0, \quad \limsup_{n \rightarrow \infty} \langle v - Pv, j(y_n - Pv) \rangle \leq 0. \quad (3.12)$$

Finally, we show that  $x_n \rightarrow Pu$ . As a matter of fact, we have

$$\begin{aligned} \|x_{n+1} - Pu\|^2 &\leq \|\beta_n(x_n - Pu) + \gamma_n(Ty_n - Pu)\|^2 + 2\alpha_n \langle u - Pu, j(x_{n+1} - Pu) \rangle \\ &\leq \beta_n \|x_n - Pu\|^2 + \gamma_n \|y_n - Pu\|^2 + 2\alpha_n \langle u - Pu, j(x_{n+1} - Pu) \rangle \\ &\leq \beta_n \|x_n - Pu\|^2 + \gamma_n \{ (1 - \delta_n) \|x_n - Pu\|^2 + 2\delta_n \langle v - Pu, j(y_n - Pu) \rangle \} \\ &\quad + 2\alpha_n \langle u - Pu, j(x_{n+1} - Pu) \rangle \\ &\leq (1 - \alpha_n) \|x_n - Pu\|^2 + 2\delta_n \langle v - Pu, j(y_n - Pu) \rangle + 2\alpha_n \langle u - Pu, j(x_{n+1} - Pu) \rangle \\ &= (1 - \alpha_n) \|x_n - Pu\|^2 + \alpha_n \lambda_n, \end{aligned} \quad (3.13)$$

where  $\lambda_n = (2\delta_n / \alpha_n) \langle v - Pu, j(y_n - Pu) \rangle + 2 \langle u - Pu, j(x_{n+1} - Pu) \rangle$ . It is easy to see from (3.12) that  $\limsup_{n \rightarrow \infty} \lambda_n \leq 0$ . Lemma 2.3 and (3.13) ensure that  $x_n \rightarrow Pu$ . This completes the proof.  $\square$

**Corollary 3.3.** Let  $C$  be a nonempty closed convex subset of a real uniformly smooth Banach space  $E$ . Let  $T : C \rightarrow C$  be a nonexpansive mapping with  $F(T) \neq \emptyset$ . Suppose that the sequences  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ , and  $\{\delta_n\}$  in  $(0, 1)$  satisfy  $\alpha_n + \beta_n + \gamma_n = 1$ ,  $n \geq 0$ . Suppose the following conditions are satisfied:

- (i)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (ii)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $\lim_{n \rightarrow \infty} \delta_n = 0$  and  $\delta_n / \alpha_n \leq M$  for some constant  $M \geq 0$ .

For fixed anchor  $u$  and arbitrary  $x_0 \in C$ , define the sequence  $\{x_n\}$  as

$$x_{n+1} = \alpha_n u + \beta_n x_n + \gamma_n T(\delta_n u + (1 - \delta_n)x_n). \quad (3.14)$$

Then, the sequence  $\{x_n\}$  converges strongly to  $Pu \in F(T)$ .

Finally, we introduce another iterative algorithm: for fixed anchor  $u \in C$  and given  $x_0 \in C$  arbitrarily, find the approximate solution  $\{x_n\}$  by the iterative algorithm:

$$\begin{aligned} x_{n+1} &= \beta_n x_n + (1 - \beta_n) A_n y_n, \\ y_n &= \alpha_n u + (1 - \alpha_n) T x_n, \quad n \geq 0, \end{aligned} \quad (3.15)$$

where  $A_n = (1/(n+1)) \sum_{i=0}^n T^i$ .

Now we prove the strong convergence of the iterative algorithm (3.15).

**Theorem 3.4.** Let  $C$  be a nonempty closed convex subset of a real uniformly smooth Banach space  $E$ . Let  $T : C \rightarrow C$  be a nonexpansive mapping with  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\}$  and  $\{\beta_n\}$  be the sequences in  $(0, 1)$ . Suppose the following conditions are satisfied:

- (i)  $\lim_{n \rightarrow \infty} \alpha_n = 0$ ;
- (ii)  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (iii)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ .

For arbitrary  $x_0 \in C$ , the sequence  $\{x_n\}$  defined by (3.15) converges strongly to  $Pu \in F(T)$ , where  $P$  is a sunny nonexpansive retraction from  $C$  onto  $F(T)$ .

*Proof.* First, we note that  $A_n : C \rightarrow C$  is nonexpansive. In fact, for any  $x, y \in C$ , we have that

$$\|A_n x - A_n y\| = \left\| \frac{1}{n+1} \sum_{i=0}^n T^i x - \frac{1}{n+1} \sum_{i=0}^n T^i y \right\| \leq \frac{1}{n+1} \sum_{i=0}^n \|T^i x - T^i y\| = \|x - y\|. \quad (3.16)$$

At the same time, for any  $x \in C$ ,

$$\|A_{n+1} x - A_n x\| = \left\| \frac{1}{n+2} \sum_{i=0}^{n+1} T^i x - \frac{1}{n+1} \sum_{i=0}^n T^i x \right\| \leq \frac{1}{(n+1)(n+2)} \sum_{i=0}^n \|T^i x\| + \frac{1}{n+2} \|T^{n+1} x\|. \quad (3.17)$$

Now we show that  $\{x_n\}$  is bounded. Indeed, for  $p \in F(T)$ , we have

$$\|y_n - p\| \leq \alpha_n \|u - p\| + (1 - \alpha_n) \|T x_n - p\| \leq \alpha_n \|u - p\| + (1 - \alpha_n) \|x_n - p\|. \quad (3.18)$$

Hence, we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq \beta_n \|x_n - p\| + (1 - \beta_n) \|A_n y_n - p\| \leq \beta_n \|x_n - p\| + (1 - \beta_n) \|y_n - p\| \\ &\leq [1 - \alpha_n(1 - \beta_n)] \|x_n - p\| + (1 - \beta_n) \alpha_n \|u - p\| \leq \max \{\|x_n - p\|, \|u - p\|\}. \end{aligned} \quad (3.19)$$

Now, an induction yields

$$\|x_n - p\| \leq \max \{\|x_0 - p\|, \|u - p\|\}. \quad (3.20)$$

Therefore,  $\{x_n\}$  is bounded, so is  $\{Tx_n\}$ .

Setting  $x_{n+1} = \beta_n x_n + (1 - \beta_n) z_n$ , where  $z_n = A_n y_n$  for all  $n \geq 0$ , then we have that

$$\begin{aligned} \|z_{n+1} - z_n\| &\leq \|A_{n+1} y_{n+1} - A_n y_{n+1}\| + \|A_n y_{n+1} - A_n y_n\| \\ &\leq \|A_{n+1} y_{n+1} - A_n y_{n+1}\| + \|y_{n+1} - y_n\|. \end{aligned} \quad (3.21)$$

Observe that

$$\begin{aligned} \|y_{n+1} - y_n\| &= \|(\alpha_{n+1} - \alpha_n)u + 1 - \alpha_{n+1}(Tx_{n+1} - Tx_n) + (\alpha_n - \alpha_{n+1})Tx_n\| \\ &\leq |\alpha_{n+1} - \alpha_n|(\|u\| + \|Tx_n\|) + (1 - \alpha_{n+1})\|x_{n+1} - x_n\|. \end{aligned} \quad (3.22)$$

So, we get

$$\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\| \leq \|A_{n+1} y_{n+1} - A_n y_{n+1}\| + |\alpha_{n+1} - \alpha_n|(\|u\| + \|Tx_n\|). \quad (3.23)$$

This together with (3.17) implies that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (3.24)$$

Consequently, from Lemma 2.2, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \|z_n - x_n\| &= 0, \\ \lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| &= \lim_{n \rightarrow \infty} (1 - \beta_n) \|x_n - z_n\| = 0. \end{aligned} \quad (3.25)$$

Next, we show that

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \quad (3.26)$$

We note from [23] that

$$\|z_n - Tz_n\| \longrightarrow 0. \quad (3.27)$$

Therefore, from (3.25), and (3.27), we have

$$\begin{aligned} \|x_n - Tx_n\| &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - z_n\| + \|z_n - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - z_n\| + \|z_n - Tz_n\| + \|Tz_n - Tx_n\| \\ &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - z_n\| + \|z_n - Tz_n\| + \|z_n - x_n\| \longrightarrow 0. \end{aligned} \quad (3.28)$$

From (3.28) and Lemma 2.5, we have

$$\limsup_{n \rightarrow \infty} \langle u - Pu, j(x_n - Pu) \rangle \leq 0. \quad (3.29)$$

Noting that

$$\|y_n - x_n\| \leq \alpha_n \|u - x_n\| + (1 - \alpha_n) \|x_n - Tx_n\| \rightarrow 0, \quad (3.30)$$

hence,

$$\limsup_{n \rightarrow \infty} \langle u - Pu, j(y_n - Pu) \rangle \leq 0. \quad (3.31)$$

Finally, applying Lemma 2.1 to (3.15), we obtain

$$\begin{aligned} \|x_{n+1} - Pu\|^2 &\leq \beta_n \|x_n - Pu\|^2 + (1 - \beta_n) \|z_n - Pu\|^2 \\ &\leq \beta_n \|x_n - Pu\|^2 + (1 - \beta_n) \|y_n - Pu\|^2 \\ &= \beta_n \|x_n - Pu\|^2 + (1 - \beta_n) \|\alpha_n(u - Pu) + (1 - \alpha_n)(Tx_n - Pu)\|^2 \\ &\leq \beta_n \|x_n - Pu\|^2 + (1 - \beta_n) [(1 - \alpha_n) \|Tx_n - Pu\|^2 + 2\alpha_n \langle u - Pu, j(y_n - Pu) \rangle] \\ &\leq \beta_n \|x_n - Pu\|^2 + (1 - \beta_n) [(1 - \alpha_n) \|x_n - Pu\|^2 + 2\alpha_n \langle u - Pu, j(y_n - Pu) \rangle] \\ &= [1 - (1 - \beta_n)\alpha_n] \|x_n - Pu\|^2 + 2(1 - \beta_n)\alpha_n \langle u - Pu, j(y_n - Pu) \rangle. \end{aligned} \quad (3.32)$$

By Lemma 2.3 and (3.32), we conclude that  $x_n \rightarrow Pu$ . This completes the proof.  $\square$

*Remark 3.5.* We drop the imposed assumptions in [21]:  $\sum_{n=1}^{\infty} |\alpha_n - \alpha_{n-1}| < \infty$  and  $\sum_{n=1}^{\infty} \delta_n < \infty$  on parameters  $\{\alpha_n\}$  and  $\{\delta_n\}$ , respectively.

## Acknowledgments

The authors are extremely grateful to the anonymous referees for their useful comments and suggestions. The first author was partially supported by Grant NSC 96-2221-E-230-003. The second author was partially supported by National Natural Science Foundation of China Grant 10771050.

## References

- [1] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, no. 2, pp. 197–228, 1967.
- [2] C. Byrne, "A unified treatment of some iterative algorithms in signal processing and image reconstruction," *Inverse Problems*, vol. 20, no. 1, pp. 103–120, 2004.
- [3] C. I. Podilchuk and R. J. Mammone, "Image recovery by convex projections using a least-squares constraint," *Journal of the Optical Society of America A*, vol. 7, no. 3, pp. 517–521, 1990.

- [4] P. L. Combettes, "The convex feasibility problem in image recovery," in *Advances in Imaging and Electron Physics*, P. Hawkes, Ed., vol. 95, pp. 155–270, Academic Press, New York, NY, USA, 1996.
- [5] P. L. Combettes, "On the numerical robustness of the parallel projection method in signal synthesis," *IEEE Signal Processing Letters*, vol. 8, no. 2, pp. 45–47, 2001.
- [6] T.-H. Kim and H.-K. Xu, "Robustness of Mann's algorithm for non-expansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 1105–1115, 2007.
- [7] H. W. Engl and A. Leitão, "A Mann iterative regularization method for elliptic Cauchy problems," *Numerical Functional Analysis and Optimization*, vol. 22, no. 7-8, pp. 861–884, 2001.
- [8] D. Youla, "Mathematical theory of image restoration by the method of convex projection," in *Image Recovery Theory and Applications*, H. Stark, Ed., pp. 29–77, Academic Press, Orlando, Fla, USA, 1987.
- [9] S. Reich, "Weak convergence theorems for non-expansive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 67, no. 2, pp. 274–276, 1979.
- [10] A. Genel and J. Lindenstrauss, "An example concerning fixed points," *Israel Journal of Mathematics*, vol. 22, no. 1, pp. 81–86, 1975.
- [11] B. Halpern, "Fixed points of nonexpanding maps," *Bulletin of the American Mathematical Society*, vol. 73, pp. 957–961, 1967.
- [12] P.-L. Lions, "Approximation de points fixes de contractions," *Comptes Rendus de l'Académie des Sciences. Séries A et B*, vol. 284, no. 21, pp. A1357–A1359, 1977.
- [13] R. Wittmann, "Approximation of fixed points of non-expansive mappings," *Archiv der Mathematik*, vol. 58, no. 5, pp. 486–491, 1992.
- [14] S. Reich, "Some problems and results in fixed point theory," in *Topological Methods in Nonlinear Functional Analysis (Toronto, Ont., 1982)*, vol. 21 of *Contemporary Mathematics*, pp. 179–187, American Mathematical Society, Providence, RI, USA, 1983.
- [15] N. Shioji and W. Takahashi, "Strong convergence of approximated sequences for non-expansive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 125, no. 12, pp. 3641–3645, 1997.
- [16] H.-K. Xu, "Another control condition in an iterative method for non-expansive mappings," *Bulletin of the Australian Mathematical Society*, vol. 65, no. 1, pp. 109–113, 2002.
- [17] H.-K. Xu, "Viscosity approximation methods for non-expansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 298, no. 1, pp. 279–291, 2004.
- [18] Y. Yao and M. A. Noor, "On viscosity iterative methods for variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 2, pp. 776–787, 2007.
- [19] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
- [20] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter non-expansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [21] Y. Su and S. Li, "Strong convergence theorems on two iterative method for non-expansive mappings," *Applied Mathematics and Computation*, vol. 181, no. 1, pp. 332–341, 2006.
- [22] Y. Liu and Z. Li, "Strong convergence theorems in reflexive Banach spaces for non-expansive mappings," *Applied Mathematics and Computation*, vol. 193, no. 2, pp. 438–445, 2007.
- [23] R. E. Bruck, "On the convex approximation property and the asymptotic behavior of nonlinear contractions in Banach spaces," *Israel Journal of Mathematics*, vol. 38, no. 4, pp. 304–314, 1981.
- [24] L.-C. Zeng and J.-C. Yao, "Implicit iteration scheme with perturbed mapping for common fixed points of a finite family of non-expansive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 11, pp. 2507–2515, 2006.
- [25] Y.-C. Liou, Y. Yao, and R. Chen, "Iteration scheme with perturbed mapping for common fixed points of a finite family of non-expansive mappings," *Fixed Point Theory and Applications*, vol. 2007, Article ID 29091, 10 pages, 2007.
- [26] Y. Yao, R. Chen, and H. Zhou, "Strong convergence to common fixed points of non-expansive mappings without commutativity assumption," *Fixed Point Theory and Applications*, vol. 2006, Article ID 89470, 8 pages, 2006.

- [27] Y.-C. Liou, Y. Yao, and K. Kimura, "Strong convergence to common fixed points of a finite family of non-expansive mappings," *Journal of Inequalities and Applications*, vol. 2007, Article ID 37513, 10 pages, 2007.
- [28] S.-S. Chang, J.-C. Yao, J. K. Kim, and L. Yang, "Iterative approximation to convex feasibility problems in Banach space," *Fixed Point Theory and Applications*, vol. 2007, Article ID 46797, 19 pages, 2007.