

Research Article

Viscosity Approximation to Common Fixed Points of Families of Nonexpansive Mappings with Weakly Contractive Mappings

A. Razani^{1,2} and S. Homaeipour¹

¹ Department of Mathematics, Faculty of Science, Imam Khomeini International University,
P.O. Box 34149-16818, Qazvin, Iran

² School of Mathematics, Institute for Research in Fundamental Sciences,
P.O. Box 19395-5746, Tehran, Iran

Correspondence should be addressed to S. Homaeipour, s.homaeipour@ikiu.ac.ir

Received 5 June 2010; Accepted 26 July 2010

Academic Editor: Brailey Sims

Copyright © 2010 A. Razani and S. Homaeipour. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let X be a reflexive Banach space which has a weakly sequentially continuous duality mapping. In this paper, we consider the following viscosity approximation sequence $x_n = \lambda_n f(x_n) + (1 - \lambda_n)T_n x_n$, where $\lambda_n \in (0, 1)$, $\{T_n\}$ is a uniformly asymptotically regular sequence, and f is a weakly contractive mapping. Strong convergence of the sequence $\{x_n\}$ is proved.

1. Introduction

Let C be a nonempty closed convex subset of a Banach space X . Recall that a self-mapping $T : C \rightarrow C$ is nonexpansive if

$$\|T(x) - T(y)\| \leq \|x - y\| \quad \forall x, y \in C. \quad (1.1)$$

Alber and Guerre-Delabriere [1] defined the weakly contractive maps in Hilbert spaces, and Rhoades [2] showed that the result of [1] is also valid in the complete metric spaces as follows.

Definition 1.1. Let (X, d) be a complete metric space. A mapping $T : X \rightarrow X$ is called weakly contractive if

$$d(Tx, Ty) \leq d(x, y) - \varphi(d(x, y)), \quad (1.2)$$

where $x, y \in X$ and $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a continuous and nondecreasing function such that $\varphi(t) = 0$ if and only if $t = 0$ and $\lim_{t \rightarrow \infty} \varphi(t) = \infty$.

Theorem 1.2. *Let $T : X \rightarrow X$ be a weakly contractive mapping, where (X, d) is a complete metric space, then T has a unique fixed point.*

In 2007, Song and Chen [3] considered the iterative sequence

$$x_n = \lambda_n f(x_n) + (1 - \lambda_n) T_n x_n, \quad n \in \{1, 2, \dots\}. \quad (1.3)$$

They proved the strong convergence of the iterative sequence $\{x_n\}$, where f is a contraction mapping and $\{T_n\}$ is a uniformly asymptotically regular sequence of nonexpansive mappings in a reflexive Banach space X , as follows.

Theorem 1.3 (see [3, Theorem 3.1]). *Let X be a reflexive Banach space which admits a weakly sequentially continuous duality mapping J from X to X^* . Suppose that C is a nonempty closed convex subset of X and $\{T_n\}, n \in \{1, 2, \dots\}$, is a uniformly asymptotically regular sequence of nonexpansive mappings from C into itself such that*

$$F := \bigcap_{n=1}^{\infty} \text{Fix}(T_n) \neq \emptyset, \quad (1.4)$$

where $\text{Fix}(T_n) := \{x \in C : x = T_n x\}, n \in \{1, 2, \dots\}$. Let $\{x_n\}$ be defined by (1.3) and $\lambda_n \in (0, 1)$, such that $\lim_{n \rightarrow \infty} \lambda_n = 0$. Then as $n \rightarrow \infty$, the sequence $\{x_n\}$ converges strongly to p , such that p is the unique solution, in F , to the variational inequality:

$$\langle f(p) - p, J(y - p) \rangle \leq 0, \quad \forall y \in F. \quad (1.5)$$

In this paper, inspired by the above results, strong convergence of sequence (1.3) is proved, where f is a weakly contractive mapping.

2. Preliminaries

A Banach space X is called strictly convex if

$$\|x\| = \|y\| = 1, \quad x \neq y \text{ implies } \frac{\|x + y\|}{2} < 1. \quad (2.1)$$

A Banach space X is called uniformly convex, if for all $\varepsilon \in [0, 2]$, there exist $\delta_\varepsilon > 0$ such that

$$\|x\| = \|y\| = 1 \quad \text{with } \|x - y\| \geq \varepsilon \text{ implies that } \frac{\|x + y\|}{2} < 1 - \delta_\varepsilon. \quad (2.2)$$

The following results are well known which can be founded in [4].

- (1) A uniformly convex Banach space X is reflexive and strictly convex.
- (2) If C is a nonempty convex subset of a strictly convex Banach space X and $T : C \rightarrow C$ is a nonexpansive mapping, then the fixed point set $F(T)$ of T is a closed convex subset of C .

By a gauge function we mean a continuous strictly increasing function φ defined on $[0, \infty)$ such that $\varphi(0) = 0$ and $\lim_{r \rightarrow \infty} \varphi(r) = \infty$. The mapping $J_\varphi : X \rightarrow 2^{X^*}$ defined by

$$J_\varphi(x) = \{x^* \in X^* : \langle x, x^* \rangle = \|x\| \|x^*\|, \|x^*\| = \varphi(\|x\|)\}, \quad \text{for each } x \in X, \quad (2.3)$$

is called the duality mapping with gauge function φ . In the case where $\varphi(t) = t$, then $J_\varphi = J$ which is the normalized duality mapping.

Proposition 2.1 (see [5]). (1) $J = I$ if and only if X is a Hilbert space.

(2) J is surjective if and only if X is reflexive.

(3) $J_\varphi(\lambda x) = \text{sign } \lambda (\varphi(|\lambda| \cdot \|x\|) / \|x\|) J(x)$ for all $x \in X \setminus \{0\}, \lambda \in \mathbb{R}$; in particular $J(-x) = -J(x)$, for all $x \in X$.

We say that a Banach space X has a weakly sequentially continuous duality mapping if there exists a gauge function φ such that the duality mapping J_φ is single-valued and continuous from the weak topology to the weak* topology of X .

We recall [6] that a Banach space X is said to satisfy Opial's condition, if for any sequence $\{x_n\}$ in X , which converges weakly to $x \in X$, we have

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\| \quad \forall y \in X, y \neq x. \quad (2.4)$$

It is known [7] that any separable Banach space can be equivalently renormed such that it satisfies Opial's condition. A space with a weakly sequentially continuous duality mapping is easily seen to satisfy Opial's condition [8].

Lemma 2.2 (see [9, Lemma 4]). *Let X be a Banach space satisfying Opial's condition and C a nonempty, closed, and convex subset of X . Suppose that $T : C \rightarrow C$ is a nonexpansive mapping. Then $I - T$ is demiclosed at zero, that is, if $\{x_n\}$ is a sequence in C which converges weakly to x and if the sequence $x_n - Tx_n$ converges strongly to zero, then $x - Tx = 0$.*

Definition 2.3 (see [3]). Let C be a nonempty closed convex subset of a Banach space X and $T_n : C \rightarrow C$, where $n \in \{1, 2, \dots\}$. Then the mapping sequence $\{T_n\}$ is called uniformly asymptotically regular on C , if for all $m \in \{1, 2, \dots\}$ and any bounded subset K of C we have

$$\lim_{n \rightarrow +\infty} \sup_{x \in K} \|T_m(T_n x) - T_n x\| = 0. \quad (2.5)$$

3. Main Result

In this section, we prove a new version of Theorem 1.3.

Theorem 3.1. *Let X be a reflexive Banach space which admits a weakly sequentially continuous duality mapping J from X to X^* . Suppose that C is a nonempty closed convex subset of X and $T_m : C \rightarrow C, m \in \{1, 2, \dots\}$, is a uniformly asymptotically regular sequence of nonexpansive mappings such that*

$$F := \bigcap_{m=1}^{\infty} \text{Fix}(T_m) \neq \emptyset. \quad (3.1)$$

Let $f : C \rightarrow C$ be a weakly contractive mapping. Suppose that $\{t_m\}$ is a sequence of positive numbers in $(0, 1)$ satisfying $\lim_{m \rightarrow \infty} t_m = 0$. Assume that $\{x_m\}$ is defined by the following iterative process:

$$x_m = t_m f(x_m) + (1 - t_m) T_m x_m, \quad m \in \{1, 2, \dots\}. \quad (3.2)$$

Then the above sequence $\{x_m\}$ converges strongly to a common fixed point p of $\{T_m\}, m \in \{1, 2, \dots\}$ such that p is the unique solution, in F , to the variational inequality

$$\langle f(p) - p, J(y - p) \rangle \leq 0, \quad \forall y \in F. \quad (3.3)$$

Proof.

Step 1. We prove the uniqueness of the solution to the variational inequality (3.3). Suppose that $p, q \in F$ are distinct solutions to (3.3). Then

$$\begin{aligned} \langle f(p) - p, J(q - p) \rangle &\leq 0, \\ \langle f(q) - q, J(p - q) \rangle &\leq 0. \end{aligned} \quad (3.4)$$

By adding up the above relations, we get

$$\begin{aligned} 0 &\geq \langle (p - f(p)) - (q - f(q)), J(p - q) \rangle \\ &\geq \langle p - q, J(p - q) \rangle - \langle f(p) - f(q), J(p - q) \rangle \\ &\geq \|p - q\|^2 - \|f(p) - f(q)\| \|J(p - q)\| \\ &\geq \|p - q\|^2 - \|p - q\|^2 + \psi(\|p - q\|) \|p - q\|. \end{aligned} \quad (3.5)$$

Thus $\psi(\|p - q\|) \|p - q\| \leq 0$, hence $p = q$. We denote by p the unique solution, in F , to (3.3).

Step 2. We show that the sequence $\{x_m\}$ is bounded. Let $q \in F$; from (3.2) we get then that

$$\begin{aligned}
\|x_m - q\|^2 &= \langle t_m(f(x_m) - q) + (1 - t_m)(T_m x_m - q), J(x_m - q) \rangle \\
&= t_m \langle (f(x_m) - f(q)) + (f(q) - q), J(x_m - q) \rangle \\
&\quad + (1 - t_m) \langle T_m x_m - T_m q, J(x_m - q) \rangle \\
&\leq t_m \|f(x_m) - f(q)\| \|J(x_m - q)\| + t_m \langle f(q) - q, J(x_m - q) \rangle \\
&\quad + (1 - t_m) \|T_m x_m - T_m q\| \|J(x_m - q)\| \\
&\leq t_m [\|x_m - q\| - \psi(\|x_m - q\|)] \|x_m - q\| + \langle f(q) - q, J(x_m - q) \rangle \\
&\quad + (1 - t_m) \|T_m x_m - T_m q\| \|J(x_m - q)\| \\
&\leq t_m [\|x_m - q\|^2 - \psi(\|x_m - q\|) \|x_m - q\| + \langle f(q) - q, J(x_m - q) \rangle] \\
&\quad + (1 - t_m) \|x_m - q\|^2 \\
&\leq \|x_m - q\|^2 - t_m \|x_m - q\| \psi(\|x_m - q\|) + t_m \|f(q) - q\| \|x_m - q\|.
\end{aligned} \tag{3.6}$$

Thus

$$\|x_m - q\| \psi(\|x_m - q\|) \leq \|f(q) - q\| \|x_m - q\|, \tag{3.7}$$

or

$$\psi(\|x_m - q\|) \leq \|f(q) - q\|. \tag{3.8}$$

Therefore $\{x_m\}$ is bounded.

Step 3. We prove that $\lim_{m \rightarrow +\infty} \|x_m - T_n x_m\| = 0$, for all $n \in \{1, 2, \dots\}$. Since the sequence $\{x_m\}$ is bounded, so $\{f(x_m)\}$ and $\{T_m x_m\}$ are bounded. Hence $\lim_{m \rightarrow \infty} t_m \|T_m x_m - f(x_m)\| = 0$, thus $\lim_{m \rightarrow \infty} \|x_m - T_m x_m\| = 0$. Let K be a bounded subset of C which contains $\{x_m\}$. Since the sequence $\{T_m\}$ is uniformly asymptotically regular, we can obtain

$$\lim_{m \rightarrow \infty} \|T_n(T_m x_m) - T_m x_m\| \leq \lim_{m \rightarrow \infty} \sup_{x \in K} \|T_n(T_m x) - T_m x\| = 0. \tag{3.9}$$

Let $m \rightarrow \infty$, then

$$\begin{aligned}
\|x_m - T_n x_m\| &\leq \|x_m - T_m x_m\| + \|T_m x_m - T_n(T_m x_m)\| + \|T_n(T_m x_m) - T_n x_m\| \\
&\leq 2\|x_m - T_m x_m\| + \|T_m x_m - T_n(T_m x_m)\| \rightarrow 0.
\end{aligned} \tag{3.10}$$

Hence $\lim_{m \rightarrow \infty} \|x_m - T_n x_m\| = 0$, for all $n \in \{1, 2, \dots\}$.

Step 4. We show that the sequence $\{x_m\}$ is sequentially compact. Since X is reflexive and $\{x_m\}$ is bounded, there exists a subsequence $\{x_{m_k}\}$ of $\{x_m\}$ such that $\{x_{m_k}\}$ is weakly convergent to $q \in C$ as $k \rightarrow \infty$. Since $\lim_{k \rightarrow \infty} \|x_{m_k} - T_n x_{m_k}\| = 0$ for all $n \in \{1, 2, \dots\}$, by Lemma 2.2, we have $q = T_n q$ for all $n \in \{1, 2, \dots\}$. Thus $q \in F$.

Step 2 implies that

$$\begin{aligned} \|x_{m_k} - q\|^2 &\leq t_{m_k} [(\|x_{m_k} - q\| - \psi(\|x_{m_k} - q\|))\|x_{m_k} - q\| + \langle f(q) - q, J(x_{m_k} - q) \rangle] \\ &\quad + (1 - t_{m_k})\|x_{m_k} - q\|^2. \end{aligned} \quad (3.11)$$

Hence

$$t_{m_k} \|x_{m_k} - q\| \psi(\|x_{m_k} - q\|) \leq t_{m_k} \langle f(q) - q, J(x_{m_k} - q) \rangle. \quad (3.12)$$

Since J is single valued and weakly sequentially continuous from X to X^* , we have

$$\limsup_{k \rightarrow \infty} \|x_{m_k} - q\| \psi(\|x_{m_k} - q\|) \leq \lim_{k \rightarrow \infty} \langle f(q) - q, J(x_{m_k} - q) \rangle = 0. \quad (3.13)$$

Thus $\lim_{k \rightarrow \infty} x_{m_k} = q$. Hence the sequence $\{x_m\}$ is sequentially compact.

Step 5. We now prove that $q \in F$ is a solution to the variational inequality (3.3). Suppose that $y \in F$, then

$$\begin{aligned} \|x_m - y\|^2 &= t_m \langle (f(x_m) - x_m) + (x_m - y), J(x_m - y) \rangle \\ &\quad + (1 - t_m) \langle T_m x_m - T_m y, J(x_m - y) \rangle \\ &\leq t_m \langle (f(x_m) - x_m), J(x_m - y) \rangle + \|x_m - y\|^2. \end{aligned} \quad (3.14)$$

Hence

$$\langle (f(x_m) - x_m), J(y - x_m) \rangle \leq 0 \quad \text{for each } m \in \{1, 2, \dots\}. \quad (3.15)$$

Since $\{x_{m_k}\} \rightarrow q$ as $k \rightarrow \infty$, we have

$$\begin{aligned} &\|(x_{m_k} - f(x_{m_k})) - (q - f(q))\| \rightarrow 0 \quad \text{as } k \rightarrow \infty, \\ &|\langle (x_{m_k} - f(x_{m_k})), J(x_{m_k} - y) \rangle - \langle (q - f(q)), J(q - y) \rangle| \\ &= |\langle (x_{m_k} - f(x_{m_k})) - (q - f(q)), J(x_{m_k} - y) \rangle + \langle (q - f(q)), J(x_{m_k} - y) - J(q - y) \rangle| \\ &\leq \|(x_{m_k} - f(x_{m_k})) - (q - f(q))\| \|x_{m_k} - y\| \\ &\quad + |\langle (q - f(q)), J(x_{m_k} - y) - J(q - y) \rangle| \rightarrow 0, \end{aligned} \quad (3.16)$$

as $k \rightarrow \infty$. Hence

$$\langle f(q) - q, J(y - q) \rangle = \lim_{k \rightarrow \infty} \langle f(x_{m_k}) - x_{m_k}, J(y - x_{m_k}) \rangle \leq 0. \quad (3.17)$$

Thus $q \in F$ is a solution to the variational inequality (3.3). By uniqueness, $q = p$. Since the sequence $\{x_m\}$ is sequentially compact and each cluster point of it is equal to p , then $\{x_m\} \rightarrow p$ as $m \rightarrow \infty$. The proof is completed. \square

It is known that [10, Example 2] in a uniformly convex Banach space E , the Cesàro means $T_n = (1/n) \sum_{j=0}^{n-1} T^j$ for nonexpansive mapping T is uniformly asymptotically regular. So we have the following corollary, which is a new version of [10, Theorem 3.2].

Corollary 3.2. *Let X be a real uniformly convex Banach space which admits a weakly sequentially continuous duality mapping J from X to X^* and C a nonempty closed convex subset of X . Suppose that $T : C \rightarrow C$ is a nonexpansive mapping, $F(T) \neq \emptyset$ and $f : C \rightarrow C$ is a weakly contractive mapping. Let $\{z_m\}$ be defined by*

$$z_m = t_m f(z_m) + (1 - t_m) \frac{1}{m+1} \sum_{j=0}^m T^j z_m, \quad m \geq 0, \quad (3.18)$$

where $t_m \in (0, 1)$ and $\lim_{m \rightarrow \infty} t_m = 0$. Then as $m \rightarrow \infty$, $\{z_m\}$ converges strongly to a fixed point p of T , where p is the unique solution in $F(T)$ to the following variational inequality:

$$\langle f(p) - p, j(u - p) \rangle \leq 0 \quad \forall u \in F(T). \quad (3.19)$$

Acknowledgment

A. Razani would like to thank the School of Mathematics of the Institute for Research in Fundamental Sciences, Teheran, Iran for supporting this paper (Grant no.89470126).

References

- [1] Ya. I. Alber and S. Guerre-Delabriere, "Principle of weakly contractive maps in Hilbert spaces," in *New Results in Operator Theory and Its Applications*, vol. 98 of *Operator Theory: Advances and Applications*, pp. 7–22, Birkhäuser, Basel, Switzerland, 1997.
- [2] B. E. Rhoades, "Some theorems on weakly contractive maps," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, pp. 2683–2693, 2001.
- [3] Y. Song and R. Chen, "Iterative approximation to common fixed points of nonexpansive mapping sequences in reflexive Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 66, no. 3, pp. 591–603, 2007.
- [4] W. Takahashi, *Nonlinear Functional Analysis. Fixed Point Theory and Its Applications*, Yokohama, Yokohama, Japan, 2000.
- [5] Z. B. Xu and G. F. Roach, "Characteristic inequalities of uniformly convex and uniformly smooth Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 157, no. 1, pp. 189–210, 1991.
- [6] Z. Opial, "Weak convergence of the sequence of successive approximations for nonexpansive mappings," *Bulletin of the American Mathematical Society*, vol. 73, pp. 591–597, 1967.
- [7] D. van Dulst, "Equivalent norms and the fixed point property for nonexpansive mappings," *Journal of the London Mathematical Society*, vol. 25, no. 1, pp. 139–144, 1982.
- [8] F. E. Browder, "Convergence theorems for sequences of nonlinear operators in Banach spaces," *Mathematische Zeitschrift*, vol. 100, pp. 201–225, 1967.

- [9] J. Górnicki, "Weak convergence theorems for asymptotically nonexpansive mappings in uniformly convex Banach spaces," *Commentationes Mathematicae Universitatis Carolinae*, vol. 30, no. 2, pp. 249–252, 1989.
- [10] Y. Song and R. Chen, "Viscosity approximate methods to Cesàro means for non-expansive mappings," *Applied Mathematics and Computation*, vol. 186, no. 2, pp. 1120–1128, 2007.