

Research Article

Monotone Generalized Nonlinear Contractions in Partially Ordered Metric Spaces

Ljubomir Ćirić,¹ Nenad Cakić,² Miloje Rajović,³ and Jeong Sheok Ume⁴

¹ Faculty of Mechanical Engineering, University of Belgrade, Kraljice Marije 16,
11 000 Belgrade, Serbia

² Faculty of Electrical Engineering, University of Belgrade, Boulevard Kralja Aleksandra 73,
11 000 Belgrade, Serbia

³ Faculty of Mechanical Engineering, University of Kragujevac, Dositejeva 19,
36 000 Kraljevo, Serbia

⁴ Department of Applied Mathematics, Changwon National University,
Changwon 641-773, South Korea

Correspondence should be addressed to Ljubomir Ćirić, lciric@rcub.bg.ac.yu

Received 29 August 2008; Accepted 9 December 2008

Recommended by Juan Jose Nieto

A concept of g -monotone mapping is introduced, and some fixed and common fixed point theorems for g -non-decreasing generalized nonlinear contractions in partially ordered complete metric spaces are proved. Presented theorems are generalizations of very recent fixed point theorems due to Agarwal et al. (2008).

Copyright © 2008 Ljubomir Ćirić et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The Banach fixed point theorem for contraction mappings has been extended in many directions (cf. [1–28]). Very recently Agarwal et al. [1] presented some new results for generalized nonlinear contractions in partially ordered metric spaces. The main idea in [1, 20, 26] involve combining the ideas of iterative technique in the contraction mapping principle with those in the monotone technique.

Recall that if (X, \leq) is a partially ordered set and $F : X \rightarrow X$ is such that for $x, y \in X$, $x \leq y$ implies $F(x) \leq F(y)$, then a mapping F is said to be non-decreasing. The main result of Agarwal et al. in [1] is the following fixed point theorem.

Theorem 1.1 (see [1, Theorem 2.2]). *Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Assume there is a non-decreasing function $\psi : [0, +\infty) \rightarrow [0, +\infty)$ with $\lim_{n \rightarrow \infty} \psi^n(t) = 0$ for each $t > 0$ and also suppose F is a non-decreasing*

mapping with

$$d(F(x), F(y)) \leq \varphi \left(\max \left\{ d(x, y), d(x, F(x)), d(y, F(y)), \frac{1}{2} [d(x, F(y)) + d(y, F(x))] \right\} \right) \quad (1.1)$$

for all $x \geq y$. Also suppose either

- (a) F is continuous or
- (b) if $\{x_n\} \subset X$ is a non-decreasing sequence with $x_n \rightarrow x$ in X , then $x_n \leq x$ for all n hold.

If there exists an $x_0 \in X$ with $x_0 \leq F(x_0)$ then F has a fixed point.

Agarwal et al. [1] observed that in certain circumstances it is possible to remove the condition that φ is non-decreasing in Theorem 1.1. So they proved the following fixed point theorem.

Theorem 1.2 (see [1, Theorem 2.3]). *Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Assume there is a continuous function $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ with $\varphi(t) < t$ for each $t > 0$ and also suppose F is a non-decreasing mapping with*

$$d(F(x), F(y)) \leq \varphi(\max \{d(x, y), d(x, F(x)), d(y, F(y))\}) \quad \forall x \geq y. \quad (1.2)$$

Also suppose either (a) or (b) holds. If there exists an $x_0 \in X$ with $x_0 \leq F(x_0)$ then F has a fixed point.

The problem to extend the result of Theorem 1.2 to mappings which satisfy (1.1) remained open. The aim of this note is to solve this problem by using more refined technique of proofs. Moreover, we introduce a concept of g -monotone mapping and prove some fixed and common fixed point theorems for g -non-decreasing generalized nonlinear contractions in partially ordered complete metric spaces.

2. Main results

Definition 2.1. Suppose (X, \leq) is a partially ordered set and $F, g : X \rightarrow X$ are mappings of X into itself. One says F is g -non-decreasing if for $x, y \in X$,

$$g(x) \leq g(y) \text{ implies } F(x) \leq F(y). \quad (2.1)$$

Now we present the main result in this paper.

Theorem 2.2. *Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Assume there is a continuous function $\varphi : [0, +\infty) \rightarrow [0, +\infty)$*

with $\varphi(t) < t$ for each $t > 0$ and also suppose $F, g : X \rightarrow X$ are such that $F(X) \subseteq g(X)$, F is a g -non-decreasing mapping and

$$d(F(x), F(y)) \leq \max \left\{ \varphi(d(g(x), g(y))), \varphi(d(g(x), F(x))), \varphi(d(g(y), F(y))), \right. \\ \left. \varphi\left(\frac{d(g(x), F(y)) + d(g(y), F(x))}{2}\right) \right\} \quad (2.2)$$

for all $x, y \in X$ for which $g(x) \geq g(y)$. Also suppose

$$\text{if } \{g(x_n)\} \subset X \text{ is a non-decreasing sequence with } g(x_n) \rightarrow g(z) \text{ in } g(X) \\ \text{then } g(x_n) \leq g(z), \quad g(z) \leq g(g(z)) \quad \forall n \text{ hold.} \quad (2.3)$$

Also suppose $g(X)$ is closed. If there exists an $x_0 \in X$ with $g(x_0) \leq F(x_0)$, then F and g have a coincidence. Further, if F, g commute at their coincidence points, then F and g have a common fixed point.

Proof. Let $x_0 \in X$ be such that $g(x_0) \leq F(x_0)$. Since $F(X) \subseteq g(X)$, we can choose $x_1 \in X$ so that $g(x_1) = F(x_0)$. Again from $F(X) \subseteq g(X)$ we can choose $x_2 \in X$ such that $g(x_2) = F(x_1)$. Continuing this process we can choose a sequence $\{x_n\}$ in X such that

$$g(x_{n+1}) = F(x_n) \quad \forall n \geq 0. \quad (2.4)$$

Since $g(x_0) \leq F(x_0)$ and $F(x_0) = g(x_1)$, we have $g(x_0) \leq g(x_1)$. Then from (2.1),

$$F(x_0) \leq F(x_1). \quad (2.5)$$

Thus, by (2.4), $g(x_1) \leq g(x_2)$. Again from (2.1),

$$F(x_1) \leq F(x_2), \quad (2.6)$$

that is, $g(x_2) \leq g(x_3)$. Continuing we obtain

$$F(x_0) \leq F(x_1) \leq F(x_2) \leq F(x_3) \leq \dots \leq F(x_n) \leq F(x_{n+1}) \leq \dots \quad (2.7)$$

In what follows we will suppose that $d(F(x_n), F(x_{n+1})) > 0$ for all n , since if $F(x_{n+1}) = F(x_n)$ for some n , then by (2.4),

$$F(x_{n+1}) = g(x_{n+1}), \quad (2.8)$$

that is, F and g have a coincidence at $x = x_{n+1}$, and so we have finished the proof. We will show that

$$d(F(x_n), F(x_{n+1})) < d(F(x_{n-1}), F(x_n)) \quad \forall n \geq 1. \quad (2.9)$$

From (2.4) and (2.7) we have that $g(x_n) \leq g(x_{n+1})$ for all $n \geq 0$. Then from (2.2) with $x = x_n$ and $y = x_{n+1}$,

$$\begin{aligned} d(F(x_n), F(x_{n+1})) \leq \max \left\{ \varphi(d(g(x_n), g(x_{n+1}))), \varphi(d(g(x_n), F(x_n))), \right. \\ \left. \varphi(d(g(x_{n+1}), F(x_{n+1}))), \right. \\ \left. \varphi\left(\frac{d(g(x_n), F(x_{n+1})) + d(g(x_{n+1}), F(x_n))}{2}\right) \right\}. \end{aligned} \quad (2.10)$$

Thus by (2.4),

$$\begin{aligned} d(F(x_n), F(x_{n+1})) \leq \max \left\{ \varphi(d(F(x_{n-1}), F(x_n))), \varphi(d(F(x_{n-1}), F(x_n))), \right. \\ \left. \varphi(d(F(x_n), F(x_{n+1}))), \varphi\left(\frac{1}{2}d(F(x_{n-1}), F(x_{n+1}))\right) \right\}. \end{aligned} \quad (2.11)$$

Hence

$$\begin{aligned} d(F(x_n), F(x_{n+1})) \leq \max \left\{ \varphi(d(F(x_{n-1}), F(x_n))), \varphi(d(F(x_n), F(x_{n+1}))), \right. \\ \left. \varphi\left(\frac{1}{2}d(F(x_{n-1}), F(x_{n+1}))\right) \right\}. \end{aligned} \quad (2.12)$$

If $d(F(x_n), F(x_{n+1})) \leq \varphi(d(F(x_{n-1}), F(x_n)))$, then (2.9) holds, as $\varphi(t) < t$ for $t > 0$.

Since we suppose that $d(F(x_n), F(x_{n+1})) > 0$ and as $\varphi(t) < t$ for $t > 0$, then $d(F(x_n), F(x_{n+1})) \leq \varphi(d(F(x_n), F(x_{n+1})))$ it is impossible.

If from (2.12) we have $d(F(x_n), F(x_{n+1})) \leq \varphi(d(F(x_{n-1}), F(x_{n+1}))/2)$, and if $d(F(x_{n-1}), F(x_{n+1}))/2 > 0$, then we have

$$\begin{aligned} d(F(x_n), F(x_{n+1})) &\leq \varphi\left(\frac{1}{2}d(F(x_{n-1}), F(x_{n+1}))\right) \\ &< \frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) \\ &\leq \frac{1}{2}d(F(x_{n-1}), F(x_n)) + \frac{1}{2}d(F(x_n), F(x_{n+1})). \end{aligned} \quad (2.13)$$

Hence

$$d(F(x_n), F(x_{n+1})) < d(F(x_{n-1}), F(x_n)). \quad (2.14)$$

Therefore, we proved that (2.9) holds.

From (2.9) it follows that the sequence $\{d(F(x_n), F(x_{n+1}))\}$ of real numbers is monotone decreasing. Therefore, there is some $\delta \geq 0$ such that

$$\lim_{n \rightarrow \infty} d(F(x_n), F(x_{n+1})) = \delta. \quad (2.15)$$

Now we will prove that $\delta = 0$. By the triangle inequality,

$$\frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) \leq \frac{1}{2}(d(F(x_{n-1}), F(x_n)) + d(F(x_n), F(x_{n+1}))). \quad (2.16)$$

Hence by (2.9),

$$\frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) < d(F(x_{n-1}), F(x_n)). \quad (2.17)$$

Taking the upper limit as $n \rightarrow \infty$ we get

$$\limsup_{n \rightarrow \infty} \frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) \leq \lim_{n \rightarrow \infty} d(F(x_{n-1}), F(x_n)). \quad (2.18)$$

If we set

$$\limsup_{n \rightarrow \infty} \frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) = b, \quad (2.19)$$

then clearly $0 \leq b \leq \delta$. Now, taking the upper limit on the both sides of (2.12) and have in mind that $\varphi(t)$ is continuous, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} d(F(x_n), F(x_{n+1})) \leq \max \left\{ \varphi \left(\lim_{n \rightarrow \infty} d(F(x_{n-1}), F(x_n)) \right), \varphi \left(\lim_{n \rightarrow \infty} d(F(x_n), F(x_{n+1})) \right), \right. \\ \left. \varphi \left(\limsup_{n \rightarrow \infty} \frac{1}{2}d(F(x_{n-1}), F(x_{n+1})) \right) \right\}. \end{aligned} \quad (2.20)$$

Hence by (2.15) and (2.19),

$$\delta \leq \max \{ \varphi(\delta), \varphi(b) \}. \quad (2.21)$$

If we suppose that $\delta > 0$, then we have

$$\delta \leq \max \{ \varphi(\delta), \varphi(b) \} < \max \{ \delta, b \} = \delta, \quad (2.22)$$

a contradiction. Thus $\delta = 0$. Therefore, we proved that

$$\lim_{n \rightarrow \infty} d(F(x_n), F(x_{n+1})) = 0. \quad (2.23)$$

Now we prove that $\{F(x_n)\}$ is a Cauchy sequence. Suppose, to the contrary, that $\{F(x_n)\}$ is not a Cauchy sequence. Then there exist an $\epsilon > 0$ and two sequences of integers $\{l(k)\}, \{m(k)\}, m(k) > l(k) \geq k$ with

$$r_k = d(F(x_{l(k)}), F(x_{m(k)})) \geq \epsilon \quad \text{for } k \in \{1, 2, \dots\}. \quad (2.24)$$

We may also assume

$$d(F(x_{l(k)}), F(x_{m(k)-1})) < \epsilon \quad (2.25)$$

by choosing $m(k)$ to be the smallest number exceeding $l(k)$ for which (2.24) holds. From (2.24), (2.25) and by the triangle inequality,

$$\epsilon \leq r_k \leq d(F(x_{l(k)}), F(x_{m(k)-1})) + d(F(x_{m(k)-1}), F(x_{m(k)})) < \epsilon + d(F(x_{m(k)-1}), F(x_{m(k)})). \quad (2.26)$$

Hence by (2.23),

$$\lim_{k \rightarrow \infty} r_k = \epsilon. \quad (2.27)$$

Since from (2.7) and (2.4) we have $g(x_{l(k)+1}) = F(x_{l(k)}) \leq F(x_{m(k)}) = g(x_{m(k)+1})$, from (2.2) and (2.4) with $x = x_{m(k)+1}$ and $y = x_{l(k)+1}$ we get

$$\begin{aligned} d(F(x_{l(k)+1}), F(x_{m(k)+1})) \leq \max \left\{ \varphi(d(F(x_{l(k)}), F(x_{m(k)}))), \varphi(d(F(x_{l(k)}), F(x_{l(k)+1}))), \right. \\ \left. \varphi(d(F(x_{m(k)}), F(x_{m(k)+1}))), \right. \\ \left. \varphi\left(\frac{d(F(x_{l(k)}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{l(k)+1}))}{2}\right) \right\}. \end{aligned} \quad (2.28)$$

Denote $\delta_n = d(F(x_n), F(x_{n+1}))$. Then we have

$$\begin{aligned} d(F(x_{l(k)+1}), F(x_{m(k)+1})) \leq \max \left\{ \varphi(r_k), \varphi(\delta_{l(k)}), \varphi(\delta_{m(k)}), \right. \\ \left. \varphi\left(\frac{d(F(x_{l(k)}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{l(k)+1}))}{2}\right) \right\}. \end{aligned} \quad (2.29)$$

Therefore, since

$$\begin{aligned} r_k &\leq d(F(x_{l(k)}), F(x_{l(k)+1})) + d(F(x_{l(k)+1}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{m(k)+1})) \\ &= \delta_{l(k)} + \delta_{m(k)} + d(F(x_{l(k)+1}), F(x_{m(k)+1})), \end{aligned} \quad (2.30)$$

we have

$$\begin{aligned} \epsilon &\leq r_k \leq \delta_{l(k)} + \delta_{m(k)} \\ &+ \max \left\{ \varphi(r_k), \varphi(\delta_{l(k)}), \varphi(\delta_{m(k)}), \varphi \left(\frac{d(F(x_{l(k)}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{l(k)+1}))}{2} \right) \right\}. \end{aligned} \quad (2.31)$$

By the triangle inequality, (2.24) and (2.25),

$$\begin{aligned} \epsilon &\leq r_k \leq d(F(x_{l(k)}), F(x_{m(k)+1})) + \delta_{m(k)}, \\ d(F(x_{l(k)}), F(x_{m(k)+1})) &\leq d(F(x_{l(k)}), F(x_{m(k)-1})) + \delta_{m(k)-1} + \delta_{m(k)} \leq \epsilon + \delta_{m(k)-1} + \delta_{m(k)}. \end{aligned} \quad (2.32)$$

From (2.32),

$$\epsilon - \delta_{m(k)} \leq d(F(x_{l(k)}), F(x_{m(k)+1})) \leq \epsilon + \delta_{m(k)-1} + \delta_{m(k)}. \quad (2.33)$$

Similarly,

$$\begin{aligned} \epsilon &\leq r_k \leq \delta_{l(k)} + d(F(x_{l(k)+1}), F(x_{m(k)})), \\ d(F(x_{l(k)+1}), F(x_{m(k)})) &\leq \delta_{l(k)} + d(F(x_{l(k)}), F(x_{m(k)-1})) + \delta_{m(k)-1} \leq \epsilon + \delta_{m(k)-1} + \delta_{m(k)}. \end{aligned} \quad (2.34)$$

Hence

$$\epsilon - \delta_{l(k)} \leq d(F(x_{m(k)}), F(x_{l(k)+1})) \leq \epsilon + \delta_{m(k)-1} + \delta_{l(k)}. \quad (2.35)$$

From (2.33) and (2.35),

$$\begin{aligned} \epsilon - \frac{\delta_{l(k)} + \delta_{m(k)}}{2} &\leq \frac{d(F(x_{l(k)}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{l(k)+1}))}{2} \\ &\leq \epsilon + \delta_{m(k)-1} + \frac{\delta_{l(k)} + \delta_{m(k)}}{2}. \end{aligned} \quad (2.36)$$

Thus from (2.36) and (2.23) we get

$$\lim_{k \rightarrow \infty} \frac{d(F(x_{l(k)}), F(x_{m(k)+1})) + d(F(x_{m(k)}), F(x_{l(k)+1}))}{2} = \epsilon. \quad (2.37)$$

Letting $n \rightarrow \infty$ in (2.31), then by (2.23), (2.27) and (2.37) we get, as φ is continuous,

$$\epsilon \leq \max \{ \varphi(\epsilon), 0, 0, \varphi(\epsilon) \} < \epsilon, \quad (2.38)$$

a contradiction. Thus our assumption (2.24) is wrong. Therefore, $\{F(x_n)\}$ is a Cauchy sequence. Since by (2.4) we have $\{F(x_n)\} = \{g(x_{n+1})\} \subseteq g(X)$ and $g(X)$ is closed, there exists $z \in X$ such that

$$\lim_{n \rightarrow \infty} g(x_n) = g(z). \quad (2.39)$$

Now we show that z is a coincidence point of F and g . Since from (2.3) and (2.39) we have $g(x_n) \leq g(z)$ for all n , then by the triangle inequality and (2.2) we get

$$\begin{aligned} d(g(z), F(z)) &\leq d(g(z), F(x_n)) + d(F(x_n), F(z)) \\ &\leq d(g(z), F(x_n)) \\ &\quad + \max \left\{ \varphi(d(g(x_n), g(z))), \varphi(d(g(x_n), F(x_n))), \right. \\ &\quad \left. \varphi(d(g(z), F(z))), \varphi\left(\frac{d(g(x_n), F(z)) + d(g(z), F(x_n))}{2}\right) \right\}. \end{aligned} \quad (2.40)$$

So letting $n \rightarrow \infty$ yields $d(g(z), F(z)) \leq \max\{\varphi(d(g(z), F(z))), \varphi(d(g(z), F(z))/2)\}$. Hence $d(g(z), F(z)) = 0$, hence $F(z) = g(z)$. Thus we proved that F and g have a coincidence.

Suppose now that F and g commute at z . Set $w = g(z) = F(z)$. Then

$$F(w) = F(g(z)) = g(F(z)) = g(w). \quad (2.41)$$

Since from (2.3) we have $g(z) \leq g(g(z)) = g(w)$ and as $g(z) = F(z)$ and $g(w) = F(w)$, from (2.2) we get

$$\begin{aligned} d(F(z), F(w)) &\leq \max \left\{ \varphi(d(g(z), g(w))), \varphi(d(g(z), F(z))), \right. \\ &\quad \left. \varphi(d(g(w), F(w))), \varphi\left(\frac{d(g(z), F(w)) + d(g(w), F(z))}{2}\right) \right\} \\ &= \varphi(d(F(z), F(w))). \end{aligned} \quad (2.42)$$

Hence $d(F(z), F(w)) = 0$, that is, $d(w, F(w)) = 0$. Therefore,

$$F(w) = g(w) = w. \quad (2.43)$$

Thus we proved that F and g have a common fixed point. \square

Remark 2.3. Note F is g -non-decreasing can be replaced by F is g -non-increasing in Theorem 2.2 provided $g(x_0) \leq F(x_0)$ is replaced by $F(x_0) \geq g(x_0)$ in Theorem 2.2.

Corollary 2.4. Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Assume there is a continuous function $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ with $\varphi(t) < t$ for each $t > 0$ and also suppose $F : X \rightarrow X$ is a non-decreasing mapping and

$$d(F(x), F(y)) \leq \max \left\{ \varphi(d(x, y)), \varphi(d(x, F(x))), \varphi(d(y, F(y))), \right. \\ \left. \varphi\left(\frac{d(x, F(y)) + d(y, F(x))}{2}\right) \right\} \quad (2.44)$$

for all $x, y \in X$ for which $x \leq y$. Also suppose either

- (i) if $\{x_n\} \subset X$ is a non-decreasing sequence with $x_n \rightarrow z$ in X then $x_n \leq z$ for all n hold or
- (ii) F is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq F(x_0)$ then F has a fixed point.

Proof. If (i) holds, then taking $g = I$ ($I =$ the identity mapping) in Theorem 2.2 we obtain Corollary 2.4. If (ii) holds, then from (2.39) with $g = I$ we get

$$z = \lim_{n \rightarrow \infty} x_{n+1} = \lim_{n \rightarrow \infty} F(x_n) = F\left(\lim_{n \rightarrow \infty} x_n\right) = F(z). \quad (2.45)$$

□

Corollary 2.5. Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Assume there is a continuous function $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ with $\varphi(t) < t$ for each $t > 0$ and also suppose $F : X \rightarrow X$ is a non-decreasing mapping and

$$d(F(x), F(y)) \leq \max \{ \varphi(d(x, y)), \varphi(d(x, F(x))), \varphi(d(y, F(y))) \} \quad (2.46)$$

for all $x, y \in X$ for which $x \leq y$. Also suppose either

- (i) if $\{x_n\} \subset X$ is a non-decreasing sequence with $x_n \rightarrow z$ in X then $x_n \leq z$ for all n hold or
- (ii) F is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq F(x_0)$ then F has a fixed point.

Remark 2.6. Since (1.2) implies (2.46) with $\varphi = \varphi$, Corollary 2.5 is a generalization of Theorem 1.2. If in addition ψ and φ are non-decreasing, then Theorem 1.2 and Corollary 2.5 are equivalent.

Taking $\varphi(t) = kt$, $0 < k < 1$, in Corollary 2.4 we obtain the following generalization of the results in [20, 26].

Corollary 2.7. Let (X, \leq) be a partially ordered set and suppose there is a metric d on X such that (X, d) is a complete metric space. Suppose $F : X \rightarrow X$ is a non-decreasing mapping and

$$d(F(x), F(y)) \leq k \max \left\{ d(x, y), d(x, F(x)), d(y, F(y)), \frac{d(x, F(y)) + d(y, F(x))}{2} \right\} \quad (2.47)$$

for all $x, y \in X$ for which $x \leq y$, where $0 < k < 1$. Also suppose either

- (i) if $\{x_n\} \subset X$ is a non-decreasing sequence with $x_n \rightarrow z$ in X then $x_n \leq z$ for all n hold or
- (ii) F is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq F(x_0)$ then F has a fixed point.

Acknowledgments

This research is financially supported by Changwon National University in 2008. The first, second, and third authors thank the Ministry of Science and Technology of Serbia for their support.

References

- [1] R. P. Agarwal, M. A. El-Gebeily, and D. O'Regan, "Generalized contractions in partially ordered metric spaces," *Applicable Analysis*, vol. 87, no. 1, pp. 109–116, 2008.
- [2] R. P. Agarwal, D. O'Regan, and M. Sambandham, "Random and deterministic fixed point theory for generalized contractive maps," *Applicable Analysis*, vol. 83, no. 7, pp. 711–725, 2004.
- [3] B. Ahmad and J. J. Nieto, "The monotone iterative technique for three-point second-order integrodifferential boundary value problems with p -Laplacian," *Boundary Value Problems*, vol. 2007, Article ID 57481, 9 pages, 2007.
- [4] D. W. Boyd and J. S. W. Wong, "On nonlinear contractions," *Proceedings of the American Mathematical Society*, vol. 20, no. 2, pp. 458–464, 1969.
- [5] A. Cabada and J. J. Nieto, "Fixed points and approximate solutions for nonlinear operator equations," *Journal of Computational and Applied Mathematics*, vol. 113, no. 1-2, pp. 17–25, 2000.
- [6] Lj. Ćirić, "Generalized contractions and fixed-point theorems," *Institut Mathématique. Publications*, vol. 12, no. 26, pp. 19–26, 1971.
- [7] Lj. Ćirić, "A generalization of Banach's contraction principle," *Proceedings of the American Mathematical Society*, vol. 45, no. 2, pp. 267–273, 1974.
- [8] Lj. Ćirić, "Fixed points of weakly contraction mappings," *Institut Mathématique. Publications*, vol. 20, no. 34, pp. 79–84, 1976.
- [9] Lj. Ćirić, "Coincidence and fixed points for maps on topological spaces," *Topology and Its Applications*, vol. 154, no. 17, pp. 3100–3106, 2007.
- [10] Lj. Ćirić and J. S. Ume, "Nonlinear quasi-contractions on metric spaces," *Praktikà tês Akadēmias Athēnon*, vol. 76, part A, pp. 132–141, 2001.
- [11] Lj. Ćirić, "Common fixed points of nonlinear contractions," *Acta Mathematica Hungarica*, vol. 80, no. 1-2, pp. 31–38, 1998.
- [12] Z. Drici, F. A. McRae, and J. Vasundhara Devi, "Fixed-point theorems in partially ordered metric spaces for operators with PPF dependence," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 2, pp. 641–647, 2007.
- [13] T. G. Bhaskar and V. Lakshmikantham, "Fixed point theorems in partially ordered metric spaces and applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 65, no. 7, pp. 1379–1393, 2006.
- [14] L. Gajić and V. Rakočević, "Quasicontraction nonself-mappings on convex metric spaces and common fixed point theorems," *Fixed Point Theory and Applications*, vol. 2005, no. 3, pp. 365–375, 2005.
- [15] N. Hussain, "Common fixed points in best approximation for Banach operator pairs with Ćirić type I -contractions," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1351–1363, 2008.
- [16] G. S. Ladde, V. Lakshmikantham, and A. S. Vatsala, *Monotone Iterative Techniques for Nonlinear Differential Equations*, Monographs, Advanced Texts and Surveys in Pure and Applied Mathematics, 27, Pitman, Boston, Mass, USA, 1985.
- [17] V. Lakshmikantham and Lj. Ćirić, "Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*. In press.
- [18] Z. Q. Liu, Z. N. Guo, S. M. Kang, and S. K. Lee, "On Ćirić type mappings with nonunique fixed and periodic points," *International Journal of Pure and Applied Mathematics*, vol. 26, no. 3, pp. 399–408, 2006.

- [19] J. J. Nieto, "An abstract monotone iterative technique," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 28, no. 12, pp. 1923–1933, 1997.
- [20] J. J. Nieto and R. Rodríguez-López, "Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations," *Order*, vol. 22, no. 3, pp. 223–239, 2005.
- [21] J. J. Nieto and R. Rodríguez-López, "Monotone method for first-order functional differential equations," *Computers & Mathematics with Applications*, vol. 52, no. 3-4, pp. 471–484, 2006.
- [22] J. J. Nieto and R. Rodríguez-López, "Existence and uniqueness of fixed point in partially ordered sets and applications to ordinary differential equations," *Acta Mathematica Sinica*, vol. 23, no. 12, pp. 2205–2212, 2007.
- [23] J. J. Nieto, R. L. Pouso, and R. Rodríguez-López, "Fixed point theorems in ordered abstract spaces," *Proceedings of the American Mathematical Society*, vol. 135, no. 8, pp. 2505–2517, 2007.
- [24] H. K. Pathak, Y. J. Cho, and S. M. Kang, "An application of fixed point theorems in best approximation theory," *International Journal of Mathematics and Mathematical Sciences*, vol. 21, no. 3, pp. 467–470, 1998.
- [25] P. Raja and S. M. Vaezpour, "Some extensions of Banach's contraction principle in complete cone metric spaces," *Fixed Point Theory and Applications*, vol. 2008, Article ID 768294, 11 pages, 2008.
- [26] A. C. M. Ran and M. C. B. Reurings, "A fixed point theorem in partially ordered sets and some applications to matrix equations," *Proceedings of the American Mathematical Society*, vol. 132, no. 5, pp. 1435–1443, 2004.
- [27] B. K. Ray, "On Ćirić's fixed point theorem," *Polska Akademia Nauk. Fundamenta Mathematicae*, vol. 94, no. 3, pp. 221–229, 1977.
- [28] D. O'Regan and A. Petrușel, "Fixed point theorems for generalized contractions in ordered metric spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 2, pp. 1241–1252, 2008.