

RESEARCH

Open Access

Modified α - ψ -contractive mappings with applications

Peyman Salimi¹, Abdul Latif^{2*} and Nawab Hussain²

*Correspondence: alatif@kau.edu.sa

²Department of Mathematics, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia
Full list of author information is available at the end of the article

Abstract

The aim of this work is to modify the notions of α -admissible and α - ψ -contractive mappings and establish new fixed point theorems for such mappings in complete metric spaces. Presented theorems provide main results of Karapinar and Samet (Abstr. Appl. Anal. 2012:793486, 2012) and Samet *et al.* (Nonlinear Anal. 75:2154-2165, 2012) as direct corollaries. Moreover, some examples and applications to integral equations are given here to illustrate the usability of the obtained results.

MSC: 46N40; 47H10; 54H25; 46T99

1 Introduction and preliminaries

Metric fixed point theory has many applications in functional analysis. The contractive conditions on underlying functions play an important role for finding solutions of metric fixed point problems. The Banach contraction principle is a remarkable result in metric fixed point theory. Over the years, it has been generalized in different directions by several mathematicians (see [1–25]). In 2012, Samet *et al.* [24] introduced the concepts of α - ψ -contractive and α -admissible mappings and established various fixed point theorems for such mappings in complete metric spaces. Afterwards Karapinar and Samet [19] generalized these notions to obtain fixed point results. The aim of this paper is to modify further the notions of α - ψ -contractive and α -admissible mappings and establish fixed point theorems for such mappings in complete metric spaces. Our results are proper generalizations of the recent results in [19, 24]. Moreover, some examples and applications to integral equations are given here to illustrate the usability of the obtained results.

Denote with Ψ the family of nondecreasing functions $\psi : [0, +\infty) \rightarrow [0, +\infty)$ such that $\sum_{n=1}^{\infty} \psi^n(t) < +\infty$ for all $t > 0$, where ψ^n is the n th iterate of ψ .

The following lemma is obvious.

Lemma 1.1 *If $\psi \in \Psi$, then $\psi(t) < t$ for all $t > 0$.*

Definition 1.1 [24] Let T be a self-mapping on a metric space (X, d) and let $\alpha : X \times X \rightarrow [0, +\infty)$ be a function. We say that T is an α -admissible mapping if

$$x, y \in X, \quad \alpha(x, y) \geq 1 \quad \implies \quad \alpha(Tx, Ty) \geq 1.$$

Definition 1.2 [24] Let T be a self-mapping on a metric space (X, d) . We say that T is an α - ψ -contractive mapping if there exist two functions $\alpha : X \times X \rightarrow [0, +\infty)$ and $\psi \in \Psi$

such that

$$\alpha(x, y)d(Tx, Ty) \leq \psi(d(x, y))$$

for all $x, y \in X$.

For the examples of α -admissible and α - ψ -contractive mappings, see [19, 24] and the examples in the next section.

2 Main results

We first modify the concept of α -admissible mapping.

Definition 2.1 Let T be a self-mapping on a metric space (X, d) and let $\alpha, \eta : X \times X \rightarrow [0, +\infty)$ be two functions. We say that T is an α -admissible mapping with respect to η if

$$x, y \in X, \quad \alpha(x, y) \geq \eta(x, y) \implies \alpha(Tx, Ty) \geq \eta(Tx, Ty).$$

Note that if we take $\eta(x, y) = 1$, then this definition reduces to Definition 1.1. Also, if we take $\alpha(x, y) = 1$, then we say that T is an η -subadmissible mapping.

Our first result is the following.

Theorem 2.1 Let (X, d) be a complete metric space and let T be an α -admissible mapping with respect to η . Assume that

$$x, y \in X, \quad \alpha(x, y) \geq \eta(x, y) \implies d(Tx, Ty) \leq \psi(M(x, y)), \tag{2.1}$$

where $\psi \in \Psi$ and

$$M(x, y) = \max \left\{ d(x, y), \frac{d(x, Tx) + d(y, Ty)}{2}, \frac{d(x, Ty) + d(y, Tx)}{2} \right\}.$$

Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq \eta(x_n, x)$ for all $n \in \mathbb{N} \cup \{0\}$.

Then T has a fixed point.

Proof Let $x_0 \in X$ be such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$. Define a sequence $\{x_n\}$ in X by $x_n = T^n x_0 = Tx_{n-1}$ for all $n \in \mathbb{N}$. If $x_{n+1} = x_n$ for some $n \in \mathbb{N}$, then $x = x_n$ is a fixed point for T and the result is proved. Hence, we suppose that $x_{n+1} \neq x_n$ for all $n \in \mathbb{N}$. Since T is a generalized α -admissible mapping with respect to η and $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$, we deduce that $\alpha(x_1, x_2) = \alpha(Tx_0, T^2x_0) \geq \eta(Tx_0, T^2x_0) = \eta(x_1, x_2)$. Continuing this process, we get $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$ for all $n \in \mathbb{N} \cup \{0\}$. Now, by (2.1) with $x = x_{n-1}$, $y = x_n$, we get

$$d(Tx_{n-1}, Tx_n) \leq \psi(M(x_{n-1}, x_n)).$$

On the other hand,

$$\begin{aligned} M(x_{n-1}, x_n) &= \max \left\{ d(x_{n-1}, x_n), \frac{d(x_{n-1}, Tx_{n-1}) + d(x_n, Tx_n)}{2}, \right. \\ &\quad \left. \frac{d(x_{n-1}, Tx_n) + d(x_n, Tx_{n-1})}{2} \right\} \\ &= \max \left\{ d(x_{n-1}, x_n), \frac{d(x_{n-1}, x_n) + d(x_n, x_{n+1})}{2}, \frac{d(x_{n-1}, x_{n+1})}{2} \right\} \\ &\leq \max \left\{ d(x_{n-1}, x_n), \frac{d(x_{n-1}, x_n) + d(x_n, x_{n+1})}{2} \right\} \\ &\leq \max \{ d(x_{n-1}, x_n), d(x_n, x_{n+1}) \}, \end{aligned}$$

which implies

$$d(x_n, x_{n+1}) \leq \psi(\max \{ d(x_{n-1}, x_n), d(x_n, x_{n+1}) \}).$$

Now, if $\max \{ d(x_{n-1}, x_n), d(x_n, x_{n+1}) \} = d(x_n, x_{n+1})$ for some $n \in \mathbb{N}$, then

$$d(x_n, x_{n+1}) \leq \psi(\max \{ d(x_{n-1}, x_n), d(x_n, x_{n+1}) \}) = \psi(d(x_n, x_{n+1})) < d(x_n, x_{n+1}),$$

which is a contradiction. Hence, for all $n \in \mathbb{N}$, we have

$$d(x_n, x_{n+1}) \leq \psi(d(x_{n-1}, x_n)).$$

By induction, we have

$$d(x_n, x_{n+1}) \leq \psi^n(d(x_0, x_1)).$$

Fix $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$\sum_{n \geq N} \psi^n(d(x_n, x_{n+1})) < \epsilon \quad \text{for all } n \in \mathbb{N}.$$

Let $m, n \in \mathbb{N}$ with $m > n \geq N$. Then, by the triangular inequality, we get

$$d(x_n, x_m) \leq \sum_{k=n}^{m-1} d(x_k, x_{k+1}) \leq \sum_{n \geq N} \psi^n(d(x_n, x_{n+1})) < \epsilon.$$

Consequently, $\lim_{m, n \rightarrow +\infty} d(x_n, x_m) = 0$. Hence $\{x_n\}$ is a Cauchy sequence. Since X is complete, there is $z \in X$ such that $x_n \rightarrow z$ as $n \rightarrow \infty$. Now, if we suppose that T is continuous, then we have

$$Tz = \lim_{n \rightarrow \infty} Tx_n = \lim_{n \rightarrow \infty} x_{n+1} = z.$$

So, z is a fixed point of T . On the other hand, since

$$\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$$

for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow z$ as $n \rightarrow \infty$, we get

$$\alpha(x_n, z) \geq \eta(x_n, z)$$

for all $n \in \mathbb{N} \cup \{0\}$. Then from (2.1) we have

$$d(x_{n+1}, Tz) \leq \psi(M(x_n, z)),$$

where

$$M(x_n, z) = \max \left\{ d(x_n, z), \frac{d(x_n, x_{n+1}) + d(z, Tz)}{2}, \frac{d(x_n, Tz) + d(z, x_{n+1})}{2} \right\}.$$

Since $M(x_n, z) > 0$, then

$$d(x_{n+1}, Tz) \leq \psi(M(x_n, z)) < M(x_n, z).$$

By taking limit as $n \rightarrow \infty$ in the above inequality, we have

$$d(z, Tz) = \lim_{n \rightarrow \infty} d(x_{n+1}, Tz) \leq \lim_{n \rightarrow \infty} M(x_n, z) = \frac{d(z, Tz)}{2},$$

which implies $d(z, Tz) = 0$, i.e., $z = Tz$. □

By taking $\eta(x, y) = 1$ in Theorem 2.1, we have the following result.

Corollary 2.1 *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that for $\psi \in \Psi$,*

$$x, y \in X, \quad \alpha(x, y) \geq 1 \quad \implies \quad d(Tx, Ty) \leq \psi(M(x, y)).$$

Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

By taking $\alpha(x, y) = 1$ in Theorem 2.1, we have the following corollary.

Corollary 2.2 *Let (X, d) be a complete metric space and let T be an η -subadmissible mapping. Assume that for $\psi \in \Psi$,*

$$x, y \in X, \quad \eta(x, y) \leq 1 \quad \implies \quad d(Tx, Ty) \leq \psi(M(x, y)).$$

Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\eta(x_0, Tx_0) \leq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\eta(x_n, x_{n+1}) \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\eta(x_n, x) \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Clearly, Corollary 2.1 implies the following results.

Corollary 2.3 (Theorem 2.1 and Theorem 2.2 of [24]) *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that for $\psi \in \Psi$,*

$$\alpha(x, y)d(Tx, Ty) \leq \psi(d(x, y))$$

holds for all $x, y \in X$. Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Corollary 2.4 (Theorem 2.3 and Theorem 2.4 of [19]) *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that for $\psi \in \Psi$,*

$$\alpha(x, y)d(Tx, Ty) \leq \psi(M(x, y)) \quad \forall x, y \in X,$$

where

$$M(x, y) = \max \left\{ d(x, y), \frac{d(x, Tx) + d(y, Ty)}{2}, \frac{d(x, Ty) + d(y, Tx)}{2} \right\}.$$

Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Example 2.1 Let $X = [0, \infty)$ be endowed with the usual metric $d(x, y) = |x - y|$ for all $x, y \in X$ and let $T : X \rightarrow X$ be defined by

$$Tx = \begin{cases} \frac{x}{2^{(x+1)}} & \text{if } x \in [0, 1], \\ \ln x + |\sin x| & \text{if } x \in (1, \infty). \end{cases}$$

Define also $\alpha : X \times X \rightarrow [0, +\infty)$ and $\psi : [0, \infty) \rightarrow [0, \infty)$ by

$$\alpha(x, y) = \begin{cases} 4 & \text{if } x, y \in [0, 1], \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad \psi(t) = \frac{1}{2}t.$$

We prove that Corollary 2.1 can be applied to T . But Theorem 2.2 of [24] and Theorem 2.4 of [19] cannot be applied to T .

Clearly, (X, d) is a complete metric space. We show that T is an α -admissible mapping. Let $x, y \in X$, if $\alpha(x, y) \geq 1$, then $x, y \in [0, 1]$. On the other hand, for all $x \in [0, 1]$ we have $Tx \leq 1$. It follows that $\alpha(Tx, Ty) \geq 1$. Hence, the assertion holds. In reason of the above arguments, $\alpha(0, T0) \geq 1$.

Now, if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, then $\{x_n\} \subset [0, 1]$ and hence $x \in [0, 1]$. This implies that $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N}$.

Let $\alpha(x, y) \geq 1$. Then $x, y \in [0, 1]$. We get

$$\begin{aligned} d(Tx, Ty) &= Ty - Tx = \frac{y}{2(y+1)} - \frac{x}{2(x+1)} \\ &= \frac{y-x}{2(1+x)(1+y)} \\ &\leq \frac{y-x}{2} = \frac{1}{2}d(x, y) \leq \frac{1}{2}M(x, y) = \psi(M(x, y)). \end{aligned}$$

That is,

$$\alpha(x, y) \geq 1 \implies d(Tx, Ty) \leq \psi(M(x, y)).$$

All of the conditions of Corollary 2.1 hold. Hence, T has a fixed point. Let $x = 0$ and $y = 1$, then

$$\alpha(0, 1)d(T0, T1) = 1 > 1/2 = \psi(d(0, 1)).$$

That is, Theorem 2.2 of [24] cannot be applied to T .

Also, by a similar method, we can show that Theorem 2.4 of [19] cannot be applied to T .

By the following simple example, we show that our results improve the results of Samet *et al.* [24] and the results of Karapinar and Samet [19].

Example 2.2 Let $X = [0, \infty)$ be endowed with the usual metric $d(x, y) = |x - y|$ for all $x, y \in X$ and let $T : X \rightarrow X$ be defined by $Tx = \frac{1}{4}x$. Also, define $\alpha : X^2 \rightarrow [0, \infty)$ by $\alpha(x, y) = 3$ and $\psi : [0, \infty) \rightarrow [0, \infty)$ by $\psi(t) = \frac{1}{2}t$.

Clearly, T is an α -admissible mapping. Also, $\alpha(x, y) = 3 \geq 1$ for all $x, y \in X$. Hence,

$$d(Tx, Ty) = \frac{1}{4}|x - y| \leq \frac{1}{2}|x - y| = \psi(d(x, y)) \leq \psi(M(x, y)).$$

Then the conditions of Corollary 2.1 hold and T has a fixed point. But if we choose $x = 4$ and $y = 8$, then

$$\alpha(4, 8)d(T4, T8) = 3 > 2 = \psi(d(4, 8)).$$

That is, Theorem 2.2 of [24] cannot be applied to T . Similarly, we can show that Theorem 2.4 of [19] cannot be applied to T . Further notice that the Banach contraction principle holds for this example.

Example 2.3 Let $X = [0, \infty)$ be endowed with the usual metric $d(x, y) = |x - y|$ for all $x, y \in X$ and let $T : X \rightarrow X$ be defined by

$$Tx = \begin{cases} \frac{1}{4}x^2 & \text{if } x \in [0, 1], \\ 2x^3 + 1 & \text{if } x \in (1, \infty). \end{cases}$$

Define also $\alpha, \eta : X \times X \rightarrow [0, +\infty)$ and $\psi : [0, \infty) \rightarrow [0, \infty)$ by

$$\eta(x, y) = \begin{cases} 1 & \text{if } x, y \in [0, 1], \\ 4 & \text{otherwise} \end{cases} \quad \text{and} \quad \psi(t) = \frac{1}{2}t.$$

We prove that Corollary 2.2 can be applied to T . But the Banach contraction principle cannot be applied to T .

Clearly, (X, d) is a complete metric space. We show that T is an η -subadmissible mapping. Let $x, y \in X$, if $\eta(x, y) \leq 1$, then $x, y \in [0, 1]$. On the other hand, for all $x \in [0, 1]$, we have $Tx \leq 1$. It follows that $\eta(Tx, Ty) \leq 1$. Also, $\eta(0, T0) \leq 1$.

Now, if $\{x_n\}$ is a sequence in X such that $\eta(x_n, x_{n+1}) \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, then $\{x_n\} \subset [0, 1]$ and hence $x \in [0, 1]$. This implies that $\eta(x_n, x) \leq 1$ for all $n \in \mathbb{N}$.

Let $\eta(x, y) \leq 1$. Then $x, y \in [0, 1]$. We get

$$d(Tx, Ty) = \frac{1}{4}|x - y||x + y| \leq \frac{1}{2}|x - y| \leq \frac{1}{2}M(x, y) = \psi(M(x, y)).$$

That is,

$$\eta(x, y) \leq 1 \implies d(Tx, Ty) \leq \psi(M(x, y)).$$

Then the conditions of Corollary 2.2 hold. Hence, T has a fixed point. Let $x = 2, y = 3$ and $r \in [0, 1)$. Then

$$d(T2, T3) = 38 > 1 > r = rd(2, 3).$$

That is, the Banach contraction principle cannot be applied to T .

From our results, we can deduce the following corollaries.

Corollary 2.5 *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that*

$$(\alpha(x, y) + \ell)^{d(Tx, Ty)} \leq (1 + \ell)^{\psi(d(x, y))} \tag{2.2}$$

holds for all $x, y \in X$, where $\psi \in \Psi$ and $\ell > 0$. Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Proof Let $\alpha(x, y) \geq 1$. Then by (2.2) we have

$$(1 + \ell)^{d(Tx, Ty)} \leq (\alpha(x, y) + \ell)^{d(Tx, Ty)} \leq (1 + \ell)^{\psi(d(x, y))}.$$

Then $d(Tx, Ty) \leq \psi(d(x, y))$. Hence, the conditions of Corollary 2.1 hold and f has a fixed point. □

Similarly, we have the following corollary.

Corollary 2.6 *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that*

$$(d(Tx, Ty) + \ell)^{\alpha(x,y)} \leq \psi(d(x, y)) + \ell \tag{2.3}$$

hold for all $x, y \in X$, where $\psi \in \Psi$ and $\ell > 0$. Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Notice that the main theorem of Dutta and Choudhury [9] remains true if ϕ is lower semi-continuous instead of continuous (see, e.g., [1, 8]).

We assume that

$$\Psi_1 = \{ \psi : [0, \infty) \rightarrow [0, \infty) \text{ such that } \psi \text{ is non-decreasing and continuous} \}$$

and

$$\Phi = \{ \varphi : [0, \infty) \rightarrow [0, \infty) \text{ such that } \varphi \text{ is lower semicontinuous} \},$$

where $\psi(t) = \varphi(t) = 0$ if and only if $t = 0$.

Theorem 2.2 *Let (X, d) be a complete metric space and let T be an α -admissible mapping with respect to η . Assume that for $\psi \in \Psi_1$ and $\varphi \in \Phi$,*

$$\begin{aligned} x, y \in X, \quad \alpha(x, Tx)\alpha(y, Ty) &\geq \eta(x, Tx)\eta(y, Ty) \\ \implies \psi(d(Tx, Ty)) &\leq \psi(d(x, y)) - \varphi(d(x, y)). \end{aligned} \tag{2.4}$$

Also, suppose that the following assertions hold:

- (i) *there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$;*
- (ii) *either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x, Tx) \geq \eta(x, Tx)$ for all $n \in \mathbb{N} \cup \{0\}$.*

Then T has a fixed point.

Proof Let $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$. Define a sequence $\{x_n\}$ in X by $x_n = T^n x_0 = Tx_{n-1}$ for all $n \in \mathbb{N}$. If $x_{n+1} = x_n$ for some $n \in \mathbb{N}$, then $x = x_n$ is a fixed point for T and the result is proved. We suppose that $x_{n+1} \neq x_n$ for all $n \in \mathbb{N}$. Since T is an α -admissible mapping with respect to η and $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$, we deduce that $\alpha(x_1, x_2) = \alpha(Tx_0, T^2x_0) \geq \eta(Tx_0, T^2x_0) = \eta(x_1, x_2)$. By continuing this process, we get $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$ for all $n \in \mathbb{N} \cup \{0\}$. Clearly,

$$\alpha(x_{n-1}, Tx_{n-1})\alpha(x_n, Tx_n) \geq \eta(x_{n-1}, Tx_{n-1})\eta(x_n, Tx_n).$$

Now, by (2.4) with $x = x_{n-1}, y = x_n$, we have

$$\psi(d(Tx_{n-1}, Tx_n)) \leq \psi(d(x_{n-1}, x_n)) - \varphi(d(x_{n-1}, x_n)),$$

which implies

$$\psi(d(x_n, x_{n+1})) \leq \psi(d(x_{n-1}, x_n)) - \varphi(d(x_{n-1}, x_n)) \leq \psi(d(x_{n-1}, x_n)). \tag{2.5}$$

Since ψ is increasing, we get

$$d(x_n, x_{n+1}) \leq d(x_{n-1}, x_n)$$

for all $n \in \mathbb{N}$. That is, $\{d_n := d(x_n, x_{n+1})\}$ is a non-increasing sequence of positive real numbers. Then there exists $r \geq 0$ such that $\lim_{n \rightarrow \infty} d_n = r$. We shall show that $r = 0$. By taking the limit infimum as $n \rightarrow \infty$ in (2.5), we have

$$\psi(r) \leq \psi(r) - \varphi(r).$$

Hence $\varphi(r) = 0$. That is, $r = 0$. Then

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \tag{2.6}$$

Suppose, to the contrary, that $\{x_n\}$ is not a Cauchy sequence. Then there is $\varepsilon > 0$ and sequences $\{m(k)\}$ and $\{n(k)\}$ such that for all positive integers k ,

$$n(k) > m(k) > k, \quad d(x_{n(k)}, x_{m(k)}) \geq \varepsilon \quad \text{and} \quad d(x_{n(k)}, x_{m(k)-1}) < \varepsilon.$$

Now, for all $k \in \mathbb{N}$, we have

$$\begin{aligned} \varepsilon &\leq d(x_{n(k)}, x_{m(k)}) \leq d(x_{n(k)}, x_{m(k)-1}) + d(x_{m(k)-1}, x_{m(k)}) \\ &< \varepsilon + d(x_{m(k)-1}, x_{m(k)}). \end{aligned}$$

Taking limit as $k \rightarrow +\infty$ in the above inequality and using (2.6), we get

$$\lim_{k \rightarrow +\infty} d(x_{n(k)}, x_{m(k)}) = \varepsilon. \tag{2.7}$$

Since

$$d(x_{n(k)}, x_{m(k)}) \leq d(x_{m(k)}, x_{m(k)+1}) + d(x_{m(k)+1}, x_{n(k)+1}) + d(x_{n(k)+1}, x_{n(k)})$$

and

$$d(x_{n(k)+1}, x_{m(k)+1}) \leq d(x_{m(k)}, x_{m(k)+1}) + d(x_{m(k)}, x_{n(k)}) + d(x_{n(k)+1}, x_{n(k)}),$$

then by taking the limit as $k \rightarrow +\infty$ in the above inequality, and by using (2.6) and (2.7), we deduce that

$$\lim_{k \rightarrow +\infty} d(x_{n(k)+1}, x_{m(k)+1}) = \varepsilon. \tag{2.8}$$

On the other hand,

$$\alpha(x_{n(k)}, Tx_{n(k)})\alpha(x_{m(k)}, Tx_{m(k)}) \geq \eta(x_{n(k)}, Tx_{n(k)})\eta(x_{m(k)}, Tx_{m(k)}).$$

Then, by (2.4) with $x = x_{n(k)}$ and $y = x_{m(k)}$, we get

$$\psi(d(x_{n(k)+1}, x_{m(k)+1})) \leq \psi(d(x_{n(k)}, x_{m(k)})) - \varphi(d(x_{n(k)}, x_{m(k)})).$$

By taking limit as $k \rightarrow \infty$ in the above inequality and applying (2.7) and (2.8), we obtain

$$\psi(\varepsilon) \leq \psi(\varepsilon) - \varphi(\varepsilon).$$

That is, $\varepsilon = 0$, which is a contradiction. Hence $\{x_n\}$ is a Cauchy sequence. Since X is complete, then there is $z \in X$ such that $x_n \rightarrow z$. First we assume that T is continuous. Then we deduce

$$Tz = \lim_{n \rightarrow \infty} Tx_n = \lim_{n \rightarrow \infty} x_{n+1} = z.$$

So, z is a fixed point of T . On the other hand, since

$$\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$$

for all $n \in \mathbb{N} \cup 0$ and $x_n \rightarrow z$ as $n \rightarrow \infty$, so

$$\alpha(z, Tz) \geq \eta(z, Tz),$$

which implies

$$\alpha(x_n, x_{n+1})\alpha(z, Tz) \geq \eta(x_n, x_{n+1})\eta(z, Tz).$$

Now, by (2.4) we get

$$\psi(d(x_{n+1}, Tz)) = \psi(d(Tx_n, Tz)) \leq \psi(d(x_n, z)) - \varphi(d(x_n, z)).$$

Passing limit inf as $n \rightarrow \infty$ in the above inequality, we have

$$\psi(d(z, Tz)) = \lim_{n \rightarrow \infty} \psi(d(x_{n+1}, Tz)) = 0.$$

That is, $z = Tz$. □

By taking $\eta(x, y) = 1$ in Theorem 2.2, we deduce the following corollary.

Corollary 2.7 *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that for $\psi \in \Psi_1$ and $\varphi \in \Phi$,*

$$x, y \in X, \quad \alpha(x, Tx)\alpha(y, Ty) \geq 1 \quad \implies \quad \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)).$$

Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x, Tx) \geq 1$.

Then T has a fixed point.

By taking $\alpha(x, y) = 1$ in Theorem 2.2, we deduce the following corollary.

Corollary 2.8 *Let (X, d) be a complete metric space and let T be an η -subadmissible mapping. Assume that for $\psi \in \Psi_1$ and $\varphi \in \Phi$,*

$$x, y \in X, \quad \eta(x, Tx)\eta(y, Ty) \leq 1 \implies \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)).$$

Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\eta(x_0, Tx_0) \leq 1$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\eta(x_n, x_{n+1}) \leq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\eta(x, Tx) \leq 1$.

Then T has a fixed point.

Example 2.4 Let $X = [0, \infty)$ be endowed with the usual metric

$$d(x, y) = \begin{cases} \max\{x, y\} & \text{if } x \neq y, \\ 0 & \text{if } x = y \end{cases}$$

for all $x, y \in X$, and let $T : X \rightarrow X$ be defined by

$$Tx = \begin{cases} \frac{x-x^2}{2} & \text{if } x \in [0, 1], \\ 2x & \text{if } x \in (1, \infty). \end{cases}$$

Define also $\alpha : X \times X \rightarrow [0, +\infty)$ and $\psi, \varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$\alpha(x, y) = \begin{cases} 1 & \text{if } x, y \in [0, 1], \\ \frac{1}{2} & \text{otherwise,} \end{cases} \quad \psi(t) = t \quad \text{and} \quad \varphi(t) = \frac{1}{2}t.$$

We prove that Corollary 2.7 can be applied to T , but the main theorem in [9] cannot be applied to T .

By a similar proof to that of Example 2.1, we show that T is an α -admissible mapping. Assume that $\alpha(x, Tx)\alpha(y, Ty) \geq 1$. Now, if $x \notin [0, 1]$, then $\alpha(x, Tx) = \frac{1}{2}$ and so $\alpha(x, Tx)\alpha(y, Ty) < 1$, which is contradiction. If $y \notin [0, 1]$. Similarly, $\alpha(x, Tx)\alpha(y, Ty) < 1$, which is contradiction. Hence, $\alpha(x, Tx)\alpha(y, Ty) \geq 1$ implies $x, y \in [0, 1]$. Therefore, we get

$$\psi(d(Tx, Ty)) = \max\left\{\frac{x-x^2}{2}, \frac{y-y^2}{2}\right\} \leq \frac{1}{2} \max\{x, y\} = \psi(d(x, y)) - \varphi(d(x, y)).$$

That is,

$$\alpha(x, Tx)\alpha(y, Ty) \geq 1 \implies \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)).$$

The conditions of Corollary 2.7 are satisfied. Hence, T has a fixed point. Let $x = 2$ and $y = 3$, then

$$\psi(d(T2, T3)) = 6 > 1/2 = \psi(d(2, 3)) - \varphi(d(2, 3)).$$

That is, the main theorem in [9] cannot be applied to T .

Example 2.5 Let $X = [0, \infty)$ be endowed with the usual metric $d(x, y) = |x - y|$ for all $x, y \in X$, and let $T : X \rightarrow X$ be defined by

$$Tx = \begin{cases} \frac{1}{5}(1 - x^3) & \text{if } x \in [0, 1], \\ |\sin(\frac{\pi}{2}x)| & \text{if } x \in (1, \infty). \end{cases}$$

Define also $\eta : X \times X \rightarrow [0, +\infty)$ and $\psi, \varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$\eta(x, y) = \begin{cases} 1 & \text{if } x, y \in [0, 1], \\ 7 & \text{otherwise,} \end{cases} \quad \psi(t) = t \quad \text{and} \quad \varphi(t) = \frac{2}{5}t.$$

We prove that Corollary 2.8 can be applied to T , but the main theorem in [9] cannot be applied to T .

By a similar proof to that of Example 2.3, we can show that T is an η -subadmissible mapping.

Assume that $\eta(x, Tx)\eta(y, Ty) \leq 1$. Now, if $x \notin [0, 1]$, then $\eta(x, Tx)\eta(y, Ty) > 1$, which is a contradiction. Similarly, $y \notin [0, 1]$ is a contradiction. Hence, $\eta(x, Tx)\eta(y, Ty) \leq 1$ implies $x, y \in [0, 1]$. We get

$$\psi(d(Tx, Ty)) = \frac{1}{5}|x - y|x^2 + xy + y^2| \leq \frac{3}{5}|x - y| = \psi(d(x, y)) - \varphi(d(x, y)).$$

That is,

$$\eta(x, Tx)\eta(y, Ty) \leq 1 \implies d(Tx, Ty) \leq \psi(d(x, y)) - \varphi(d(x, y)).$$

Then the conditions of Corollary 2.8 hold and T has a fixed point. Let $x = 2, y = 3$. Then $T2 = 0$ and $T3 = 1$, which implies

$$\psi(d(T2, T3)) = 1 > \frac{3}{5} = \psi(d(2, 3)) - \varphi(d(2, 3)).$$

That is, the main theorem in [9] cannot be applied to T .

In 1984 Khan *et al.* [20] proved the following theorem.

Theorem 2.3 Let (X, d) be a complete metric space and let T be a self-mapping on X . Assume that

$$\psi(d(Tx, Ty)) \leq c\psi(d(x, y)) \quad \forall x, y \in X,$$

where $\psi \in \Psi_1$ and $0 < c < 1$. Then T has a unique fixed point.

Theorem 2.4 *Let (X, d) be a complete metric space and let T be a generalized α -admissible mapping with respect to η . Assume that*

$$x, y \in X, \quad \alpha(x, x)\alpha(y, y) \geq \eta(x, x)\eta(y, y) \implies \psi(d(Tx, Ty)) \leq c\psi(d(x, y)), \quad (2.9)$$

where $\psi \in \Psi_1$ and $0 < c < 1$. Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\alpha(x_0, x_0) \geq \eta(x_0, x_0)$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_n) \geq \eta(x_n, x_n)$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x, x) \geq \eta(x, x)$.

Then T has a fixed point.

Proof Let $x_0 \in X$ such that $\alpha(x_0, x_0) \geq \eta(x_0, x_0)$. Define a sequence $\{x_n\}$ in X by $x_n = T^n x_0 = Tx_{n-1}$ for all $n \in \mathbb{N}$. If $x_{n+1} = x_n$ for some $n \in \mathbb{N}$, then $x = x_n$ is a fixed point for T and the result is proved. Hence, we suppose that $x_{n+1} \neq x_n$ for all $n \in \mathbb{N}$. Since T is a generalized α -admissible mapping with respect to η and $\alpha(x_0, x_0) \geq \eta(x_0, x_0)$, we deduce that $\alpha(x_1, x_1) = \alpha(Tx_0, Tx_0) \geq \eta(Tx_0, Tx_0) = \eta(x_1, x_1)$. By continuing this process, we get $\alpha(x_n, x_n) \geq \eta(x_n, x_n)$ for all $n \in \mathbb{N} \cup \{0\}$. Clearly,

$$\alpha(x_{n-1}, x_{n-1})\alpha(x_n, x_n) \geq \eta(x_{n-1}, x_{n-1})\eta(x_n, x_n).$$

Now, by (2.9) with $x = x_{n-1}$, $y = x_n$, we have

$$\psi(d(x_n, x_{n+1})) = \psi(d(Tx_{n-1}, Tx_n)) \leq c\psi(d(x_{n-1}, x_n)) < \psi(d(x_{n-1}, x_n)). \quad (2.10)$$

Since, ψ is increasing, we get

$$d(x_n, x_{n+1}) \leq d(x_{n-1}, x_n)$$

for all $n \in \mathbb{N}$. That is, $\{d_n := d(x_n, x_{n+1})\}$ is a non-increasing sequence of positive real numbers. Then there exists $r \geq 0$ such that $\lim_{n \rightarrow \infty} d_n = r$. We shall show that $r = 0$. By taking the limit as $n \rightarrow \infty$ in (2.10), we have

$$\psi(r) \leq c\psi(r),$$

which implies $\psi(r) = 0$, i.e., $r = 0$. Then

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \quad (2.11)$$

Suppose, to the contrary, that $\{x_n\}$ is not a Cauchy sequence. Proceeding as in the proof of Theorem 2.2, there exists $\epsilon > 0$ such that for all $k \in \mathbb{N}$ there exist $n(k), m(k) \in \mathbb{N}$ with $m(k) > n(k) \geq k$ such that

$$\lim_{k \rightarrow +\infty} d(x_{n(k)}, x_{m(k)}) = \epsilon \quad (2.12)$$

and

$$\lim_{k \rightarrow +\infty} d(x_{n(k)+1}, x_{m(k)+1}) = \epsilon. \quad (2.13)$$

Clearly,

$$\alpha(x_{n(k)}, x_{n(k)})\alpha(x_{m(k)}, x_{m(k)}) \geq \eta(x_{n(k)}, x_{n(k)})\eta(x_{m(k)}, x_{m(k)}).$$

Then, by (2.9) with $x = x_{n(k)}$ and $y = x_{m(k)}$, we get

$$\psi(d(x_{n(k)+1}, x_{m(k)+1})) = \psi(d(Tx_{n(k)}, Tx_{m(k)})) \leq c\psi(d(x_{n(k)}, x_{m(k)})).$$

Taking limit as $k \rightarrow \infty$ in the above inequality and applying (2.12) and (2.13), we get

$$\psi(\epsilon) \leq c\psi(\epsilon),$$

and so $\epsilon = 0$, which is a contradiction. Hence $\{x_n\}$ is a Cauchy sequence. Since X is complete, then there is $z \in X$ such that $x_n \rightarrow z$. First, we assume that T is continuous. Then, we deduce

$$Tz = \lim_{n \rightarrow \infty} Tx_n = \lim_{n \rightarrow \infty} x_{n+1} = z.$$

So, z is a fixed point of T . On the other hand, since

$$\alpha(x_n, x_n) \geq \eta(x_n, x_n)$$

for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow z$ as $n \rightarrow \infty$, we get

$$\alpha(z, z) \geq \eta(z, z),$$

which implies

$$\alpha(z, z)\alpha(x_n, x_n) \geq \eta(z, z)\eta(x_n, x_n).$$

Then by (2.12) we deduce

$$\psi(d(x_{n+1}, Tz)) = \psi(d(Tx_n, Tz)) \leq c\psi(d(x_n, z)).$$

Taking limit as $n \rightarrow \infty$ in the above inequality, we have

$$\psi(d(z, Tz)) \leq \psi(0) = 0$$

and then $z = Tz$. □

Corollary 2.9 *Let (X, d) be a complete metric space and let T be an α -admissible mapping. Assume that*

$$x, y \in X, \quad \alpha(x, x)\alpha(y, y) \geq 1 \quad \implies \quad \psi(d(Tx, Ty)) \leq c\psi(d(x, y)),$$

where $\psi \in \Psi_1$ and $0 < c < 1$. Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\alpha(x_0, x_0) \geq 1$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\alpha(x_n, x_n) \geq 1$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\alpha(x, x) \geq 1$.

Then T has a fixed point.

Corollary 2.10 Let (X, d) be a complete metric space and let T be a generalized α -admissible mapping with respect to η . Assume that

$$x, y \in X, \quad \eta(x, x)\eta(y, y) \leq 1 \implies \psi(d(Tx, Ty)) \leq c\psi(d(x, y)),$$

where $\psi \in \Psi_1$ and $0 < c < 1$. Also, suppose that the following assertions hold:

- (i) there exists $x_0 \in X$ such that $\eta(x_0, x_0) \leq 1$;
- (ii) either T is continuous or for any sequence $\{x_n\}$ in X with $\eta(x_n, x_n) \leq 1$ for all $n \in \mathbb{N} \cup 0$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, we have $\eta(x, x) \leq 1$.

Then T has a fixed point.

Example 2.6 Let $X = [0, \infty)$ be endowed with the usual metric

$$d(x, y) = \begin{cases} \max\{x, y\} & \text{if } x \neq y, \\ 0 & \text{if } x = y \end{cases}$$

for all $x, y \in X$, and let $T : X \rightarrow X$ be defined by

$$Tx = \begin{cases} \frac{x^3 - x^5}{8} & \text{if } x \in [0, 1], \\ 2x^2 + |(x - 2)(x - 3)| & \text{if } x \in (1, \infty). \end{cases}$$

Define also $\alpha : X \times X \rightarrow [0, +\infty)$ and $\psi, \varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$\alpha(x, y) = \begin{cases} 1 & \text{if } x, y \in [0, 1], \\ \frac{1}{2} & \text{otherwise} \end{cases} \quad \text{and} \quad \psi(t) = t^2.$$

We prove that Corollary 2.9 can be applied to T . But Theorem 2.3 cannot be applied to T .

By a similar proof to that of Example 2.1, we show that T is an α -admissible mapping. Assume that $\alpha(x, x)\alpha(y, y) \geq 1$. Now, if $x \notin [0, 1]$, then $\alpha(x, x) = \frac{1}{2}$ and so $\alpha(x, x)\alpha(y, y) < 1$, which is contradiction. If $y \notin [0, 1]$. Similarly, $\alpha(x, x)\alpha(y, y) < 1$, which is contradiction. Hence, $\alpha(x, x)\alpha(y, y) \geq 1$ implies $x, y \in [0, 1]$. Therefore, we get

$$\psi(d(Tx, Ty)) = \left(\max \left\{ \frac{x^3 - x^5}{8}, \frac{y^3 - y^5}{8} \right\} \right)^2 \leq \frac{1}{16} (\max\{x, y\})^2 = \frac{1}{16} \psi(d(x, y)).$$

That is,

$$\alpha(x, x)\alpha(y, y) \geq 1 \implies \psi(d(Tx, Ty)) \leq \frac{1}{16} \psi(d(x, y)).$$

Then the conditions of Corollary 2.9 hold. Hence, T has a fixed point. Let $x = 2$ and $y = 3$, then $T2 = 8$ and $T3 = 18$, and hence

$$\psi(d(T2, T3)) = 100 > \frac{1}{16} = \frac{1}{16}\psi(d(2, 3)).$$

That is, Theorem 2.3 cannot be applied to T .

3 Application to the existence of solutions of integral equations

Integral equations like (3.1) were studied in many papers (see [2, 11] and references therein). In this section, we look for a nonnegative solution to (3.1) in $X = C([0, T], \mathbb{R})$. Let $X = C([0, T], \mathbb{R})$ be the set of real continuous functions defined on $[0, T]$ and let $d : X \times X \rightarrow \mathbb{R}_+$ be defined by

$$d(x, y) = \|x - y\|_\infty$$

for all $x, y \in X$. Then (X, d) is a complete metric space.

Consider the integral equation

$$x(t) = p(t) + \int_0^T S(t, s)f(s, x(s)) ds, \tag{3.1}$$

and let $F : X \rightarrow X$ defined by

$$F(x)(t) = p(t) + \int_0^T S(t, s)f(s, x(s)) ds. \tag{3.2}$$

We assume that

- (A) $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous;
- (B) $p : [0, T] \rightarrow \mathbb{R}$ is continuous;
- (C) $S : [0, T] \times \mathbb{R} \rightarrow [0, +\infty)$ is continuous;
- (D) there exist $\psi \in \Psi$ and $\theta : X \times X \rightarrow \mathbb{R}$ such that if $\theta(x, y) \geq 0$ for $x, y \in X$, then for every $s \in [0, T]$ we have

$$\begin{aligned} &0 \leq f(s, x(s)) - f(s, y(s)) \\ &\leq \psi \left(\max \left\{ |x(s) - y(s)|, \frac{1}{2} [|x(s) - F(x(s))| + |y(s) - F(y(s))|], \right. \right. \\ &\quad \left. \left. \frac{1}{2} [|x(s) - F(y(s))| + |y(s) - F(x(s))|] \right\} \right); \end{aligned}$$

- (F) there exists $x_0 \in X$ such that $\theta(x_0, F(x_0)) \geq 0$;
- (G) if $\theta(x, y) \geq 0, x, y \in X$, then $\theta(Fx, Fy) \geq 0$;
- (H) if $\{x_n\}$ is a sequence in X such that $\theta(x_n, x_{n+1}) \geq 0$ for all $n \in \mathbb{N} \cup \{0\}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, then $\theta(x_n, x) \geq 0$ for all $n \in \mathbb{N} \cup \{0\}$;
- (J) $\int_0^T S(t, s) ds \leq 1$ for all $t \in [0, T]$ and $s \in \mathbb{R}$.

Theorem 3.1 *Under assumptions (A)-(J), the integral equation (3.1) has a solution in $X = C([0, T], \mathbb{R})$.*

Proof Consider the mapping $F : X \rightarrow X$ defined by (3.2).

By the condition (D), we deduce

$$\begin{aligned}
 & |F(x)(t) - F(y)(t)| \\
 &= \left| \int_0^T S(t,s)[f(s,x(s)) - f(s,y(s))] ds \right| \\
 &\leq \int_0^T S(t,s)|f(s,x(s)) - f(s,y(s))| ds \\
 &\leq \int_0^T S(t,s) \left[\psi \left(\max \left\{ |x(s) - y(s)|, \frac{1}{2} [|x(s) - F(x(s))| + |y(s) - F(y(s))|] \right\}, \right. \right. \\
 &\quad \left. \left. \frac{1}{2} [|x(s) - F(y(s))| + |y(s) - F(x(s))|] \right) \right] ds \\
 &\leq \int_0^T S(t,s) \left[\psi \left(\max \left\{ \|x(s) - y(s)\|, \frac{1}{2} [\|x(s) - F(x(s))\| + \|y(s) - F(y(s))\|] \right\}, \right. \right. \\
 &\quad \left. \left. \frac{1}{2} [\|x(s) - F(y(s))\| + \|y(s) - F(x(s))\|] \right) \right] ds \\
 &= \left(\int_0^T S(t,s) ds \right) \psi \left(\max \left\{ \|x(s) - y(s)\|, \frac{1}{2} [\|x(s) - F(x(s))\| + \|y(s) - F(y(s))\|] \right\}, \right. \\
 &\quad \left. \frac{1}{2} [\|x(s) - F(y(s))\| + \|y(s) - F(x(s))\|] \right) \\
 &\leq \psi \left(\max \left\{ \|x(s) - y(s)\|, \frac{1}{2} [\|x(s) - F(x(s))\| + \|y(s) - F(y(s))\|] \right\}, \right. \\
 &\quad \left. \frac{1}{2} [\|x(s) - F(y(s))\| + \|y(s) - F(x(s))\|] \right).
 \end{aligned}$$

Then

$$\begin{aligned}
 \|Fx - Fy\|_\infty &\leq \psi \left(\max \left\{ \|x(s) - y(s)\|, \frac{1}{2} [\|x(s) - F(x(s))\| + \|y(s) - F(y(s))\|] \right\}, \right. \\
 &\quad \left. \frac{1}{2} [\|x(s) - F(y(s))\| + \|y(s) - F(x(s))\|] \right).
 \end{aligned}$$

Now, define $\alpha : X \times X \rightarrow [0, +\infty)$ by

$$\alpha(x,y) = \begin{cases} 1 & \text{if } \theta(x,y) \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

That is, $\alpha(x,y) \geq 1$ implies

$$\begin{aligned}
 \|Fx - Fy\|_\infty &\leq \psi \left(\max \left\{ \|x(s) - y(s)\|, \frac{1}{2} [\|x(s) - F(x(s))\| + \|y(s) - F(y(s))\|] \right\}, \right. \\
 &\quad \left. \frac{1}{2} [\|x(s) - F(y(s))\| + \|y(s) - F(x(s))\|] \right), \\
 \|Fx - Fy\|_\infty &\leq \psi \left(\max \left\{ \|x - y\|_\infty, \frac{1}{2} [\|x - F(x)\|_\infty + \|y - F(y)\|_\infty] \right\}, \right. \\
 &\quad \left. \frac{1}{2} [\|x - F(y)\|_\infty + \|y - F(x)\|_\infty] \right).
 \end{aligned}$$

All of the hypotheses of Corollary 2.1 are satisfied, and hence the mapping F has a fixed point that is a solution in $X = C([0, T], \mathbb{R})$ of the integral equation (3.1). \square

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Astara Branch, Islamic Azad University, Astara, Iran. ²Department of Mathematics, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia.

Acknowledgements

The authors thank the referees for their valuable comments and suggestions. This research was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah. Therefore, the second and third authors acknowledge with thanks DSR, KAU for financial support. The first author is thankful for support of Astara Branch, Islamic Azad University, during this research.

Received: 14 February 2013 Accepted: 21 May 2013 Published: 10 June 2013

References

1. Abbas, M, Dorić, D: Common fixed point theorem for four mappings satisfying generalized weak contractive condition. *Filomat* **24**(2), 1-10 (2010)
2. Agarwal, RP, Hussain, N, Taoudi, M-A: Fixed point theorems in ordered Banach spaces and applications to nonlinear integral equations. *Abstr. Appl. Anal.* **2012**, Article ID 245872 (2012)
3. Akbar, F, Khan, AR: Common fixed point and approximation results for noncommuting maps on locally convex spaces. *Fixed Point Theory Appl.* **2009**, Article ID 207503 (2009)
4. Berinde, V, Vetro, F: Common fixed points of mappings satisfying implicit contractive conditions. *Fixed Point Theory Appl.* **2012**, 105 (2012)
5. Ćirić, LB: A generalization of Banach's contraction principle. *Proc. Am. Math. Soc.* **45**, 267-273 (1974)
6. Ćirić, L, Hussain, N, Cakic, N: Common fixed points for Ćirić type f -weak contraction with applications. *Publ. Math. (Debr.)* **76**(1-2), 31-49 (2010)
7. Ćirić, L, Abbas, M, Saadati, R, Hussain, N: Common fixed points of almost generalized contractive mappings in ordered metric spaces. *Appl. Math. Comput.* **217**, 5784-5789 (2011)
8. Dorić, D: Common fixed point for generalized (ψ, ϕ) -weak contractions. *Appl. Math. Lett.* **22**, 1896-1900 (2009)
9. Dutta, PN, Choudhury, BS: A generalization of contraction principle in metric spaces. *Fixed Point Theory Appl.* **2008**, Article ID 406368 (2008)
10. Hussain, N, Berinde, V, Shafiqat, N: Common fixed point and approximation results for generalized ϕ -contractions. *Fixed Point Theory* **10**, 111-124 (2009)
11. Hussain, N, Khan, AR, Agarwal, RP: Krasnosel'skii and Ky Fan type fixed point theorems in ordered Banach spaces. *J. Nonlinear Convex Anal.* **11**(3), 475-489 (2010)
12. Hussain, N, Khamsi, MA, Latif, A: Common fixed points for JH -operators and occasionally weakly biased pairs under relaxed conditions. *Nonlinear Anal.* **74**, 2133-2140 (2011)
13. Hussain, N, Kadelburg, Z, Radenovic, S, Al-Solamy, FR: Comparison functions and fixed point results in partial metric spaces. *Abstr. Appl. Anal.* **2012**, Article ID 605781 (2012)
14. Hussain, N, Dorić, D, Kadelburg, Z, Radenović, S: Suzuki-type fixed point results in metric type spaces. *Fixed Point Theory Appl.* **2012**, 126 (2012)
15. Hussain, N, Karapinar, E, Salimi, P, Akbar, F: α -Admissible mappings and related fixed point theorems. *J. Inequal. Appl.* **2013**, 114 (2013)
16. Hussain, N, Karapinar, E, Salimi, P, Vetro, P: Fixed point results for G^m -Meir-Keeler contractive and $G(\alpha, \psi)$ -Meir-Keeler contractive mappings. *Fixed Point Theory Appl.* **2013**, 34 (2013)
17. Jachymski, J: Equivalent conditions and the Meir-Keeler type theorems. *J. Math. Anal. Appl.* **194**(1), 293-303 (1995)
18. Kadelburg, Z, Radenović, S: Meir-Keeler-type conditions in abstract metric spaces. *Appl. Math. Lett.* **24**(8), 1411-1414 (2011)
19. Karapinar, E, Samet, B: Generalized α - ψ contractive type mappings and related fixed point theorems with applications. *Abstr. Appl. Anal.* **2012**, Article ID 793486 (2012)
20. Khan, MS, Swaleh, M, Sessa, S: Fixed point theorems by altering distances between the points. *Bull. Aust. Math. Soc.* **30**, 1-9 (1984)
21. Latif, A, Albar, WA: Fixed point results in complete metric spaces. *Demonstr. Math.* **XLI**(1), 145-150 (2008)
22. Latif, A, Abdou, AAN: Multivalued generalized nonlinear contractive maps and fixed points. *Nonlinear Anal.* **74**, 1436-1444 (2011)
23. Latif, A, Al-Mezel, SA: Fixed point results in quasimetric spaces. *Fixed Point Theory Appl.* **2011**, Article ID 178306 (2011)
24. Samet, B, Vetro, C, Vetro, P: Fixed point theorem for α - ψ contractive type mappings. *Nonlinear Anal.* **75**, 2154-2165 (2012)
25. Suzuki, T: A generalized Banach contraction principle that characterizes metric completeness. *Proc. Am. Math. Soc.* **136**, 1861-1869 (2008)

doi:10.1186/1687-1812-2013-151

Cite this article as: Salimi et al.: Modified α - ψ -contractive mappings with applications. *Fixed Point Theory and Applications* 2013 **2013**:151.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
