

RESEARCH

Open Access

Convergence theorems for a system of equilibrium problems and fixed point problems of a strongly nonexpansive sequence

Atid Kangtunyakarn*

*Correspondence:
beawrock@hotmail.com
Department of Mathematics,
Faculty of Science, King Mongkut's
Institute of Technology Ladkrabang,
Bangkok 10520, Thailand

Abstract

The purpose of this paper is to prove a strong convergence theorem of an iterative scheme associated to a strongly nonexpansive sequence for finding a common element of the set of equilibrium problems and the set of fixed point problems of a pair of sequences of nonexpansive mappings where one of them is a strongly nonexpansive sequence. Moreover, in the last section, by using our main result, we obtain a strong convergence theorem of an iterative scheme associated to a strongly nonexpansive sequence for finding a common element of the set of a finite family of equilibrium problems and the set of fixed point problems of a pair of sequences of nonexpansive mappings where one of them is a strongly nonexpansive sequence in a Hilbert space, and we also give some examples to support our main result.

Keywords: nonexpansive mappings; strongly nonexpansive sequence; equilibrium problem; fixed point

1 Introduction

Throughout this paper, we assume that H is a real Hilbert space with the inner product $\langle \cdot, \cdot \rangle$ and the norm $\| \cdot \|$. A mapping T of C into itself is called *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in H$. The set of fixed points of T is denoted by $F(T)$, i.e., $F(T) = \{x \in H : Tx = x\}$. It is known that $F(T)$ is closed and convex if T is nonexpansive. Let P_C be a metric projection of H onto C , i.e., for $x \in H$, $P_C x$ satisfies the property

$$\|x - P_C x\| = \min_{y \in C} \|x - y\|.$$

We use " \rightharpoonup " and " \rightarrow " to denote weak and strong convergence, respectively. Let $\{T_n\}$ be a sequence of mappings of C into H . The set of common fixed points of $\{T_n\}$ is denoted by $F(\{T_n\}) = \bigcap_{n=1}^{\infty} F(T_n)$. Recall the main concepts as follows:

- (1) A sequence $\{z_n\}$ in C is said to be an *approximate fixed point sequence* of $\{T_n\}$ if $z_n - T_n z_n \rightarrow 0$. The set of all *bounded* approximate fixed point sequences of $\{T_n\}$ is denoted by $\tilde{F}(\{T_n\})$; see [1]. It is clear that if $\{T_n\}$ has a common fixed point, then $\tilde{F}(\{T_n\})$ is nonempty.

- (2) A sequence $\{T_n\}$ is said to be a *strongly nonexpansive sequence* if each T_n is nonexpansive and

$$x_n - y_n - (T_n x_n - T_n y_n) \rightarrow 0,$$

whenever $\{x_n\}$ and $\{y_n\}$ are sequences in C such that $\{x_n - y_n\}$ is bounded and $\|x_n - y_n\| - \|T_n x_n - T_n y_n\| \rightarrow 0$.

- (3) A sequence $\{T_n\}$ having a common fixed point is said to satisfy the *condition (Z)* if every weak cluster point of $\{x_n\}$ is a common fixed point whenever $\{x_n\} \in \tilde{F}(\{T_n\})$.
(4) A sequence $\{T_n\}$ of nonexpansive mappings of C into H is said to satisfy the *condition (R)* if

$$\lim_{n \rightarrow \infty} \sup_{y \in D} \|T_{n+1}y - T_n y\| = 0$$

for every nonempty bounded subset D of C ; see [2].

Example 1.1 Let \mathbb{R} be a set of real numbers. For every $n \in \mathbb{N}$, the mapping $T_n : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $T_n x = \frac{1}{n}x$ for all $x \in \mathbb{R}$.

Then $\{T_n\}$ is a nonexpansive sequence, but it is not a strongly nonexpansive sequence.

Example 1.2 For every $n \in \mathbb{N}$, the mapping $T_n : [0, 1] \rightarrow [0, 1]$ is defined by $T_n x = (1 - \frac{1}{n})x$ for all $x \in [0, 1]$.

Then $\{T_n\}$ is a strongly nonexpansive sequence.

Solution It is easy to see that T_n is a nonexpansive mapping for all $n \in \mathbb{N}$.

Let $\{x_n\}$ and $\{y_n\}$ be sequences in $[0, 1]$ with $\{x_n - y_n\}$ being bounded and $|x_n - y_n| - |T_n x_n - T_n y_n| \rightarrow 0$ as $n \rightarrow \infty$.

Since $x_n - y_n - (T_n x_n - T_n y_n) = \frac{1}{n}(x_n - y_n)$, for all $n \in \mathbb{N}$, then we have

$$x_n - y_n - (T_n x_n - T_n y_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then $\{T_n\}$ is a strongly nonexpansive sequence.

Let $G : C \times C \rightarrow \mathbb{R}$ be a bifunction. The equilibrium problem for G is to determine its equilibrium points, *i.e.*, the set

$$EP(G) = \{x \in C : G(x, y) \geq 0, \forall y \in C\}.$$

It is a unified model of several problems, namely, variational inequality problem, complementary problem, saddle point problem, optimization problem, fixed point problem, *etc.*; see [3–5]. Several iterative methods have been proposed to solve the equilibrium problem; see, for instance, [6–8]. In 2005, Combettes and Hirstoaga [4] introduced some iterative schemes of finding the best approximation to the initial data when $EP(G)$ is nonempty and proved a strong convergence theorem.

Also in [4], Combettes and Hiratoaga, following [3], defined $S_r : H \rightarrow C$ by

$$S_r(x) = \left\{ z \in C : G(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}.$$

They proved that under suitable hypotheses S_r is single-valued and firmly nonexpansive with $F(S_r) = EP(G)$.

In 2007, Takahashi and Takahashi [9] proved the following theorem.

Theorem 1.3 *Let C be a nonempty closed convex subset of H . Let G be a bifunction from $C \times C$ to \mathbb{R} satisfying*

- (A1) $G(x, x) = 0, \forall x \in C$;
- (A2) G is monotone, i.e., $G(x, y) + G(y, x) \leq 0, \forall x, y \in C$;
- (A3) $\forall x, y, z \in C, \lim_{t \rightarrow 0^+} G(tz + (1-t)x, y) \leq G(x, y)$;
- (A4) $\forall x \in C, y \mapsto G(x, y)$ is convex and lower semicontinuous;

and let S be a nonexpansive mapping of C into H such that $F(S) \cap EP(G) \neq \emptyset$. Let f be a contraction of H into itself, and let $\{x_n\}$ and $\{u_n\}$ be sequences generated by $x_1 \in H$ and

$$G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C,$$

$$x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n) S u_n$$

for all $n \in \mathbb{N}$, where $\{\alpha_n\} \subset [0, 1]$ and $\{r_n\} \subset (0, 1)$ satisfy

- (C1) $\lim_{n \rightarrow \infty} \alpha_n = 0$;
- (C2) $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (C3) $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$;

and $\liminf_{n \rightarrow \infty} r_n > 0$ and $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$.

Then $\{x_n\}$ and $\{u_n\}$ converge strongly to $z \in F(S) \cap EP(G)$, where $z = P_{F(S) \cap EP(G)} f(z)$.

Very recently, in 2011, Aoyama and Kimura [10] proved a strong convergence theorem of the iterative scheme of $\{x_n\}$ associated to a strongly nonexpansive sequence as follows.

Theorem 1.4 *Let H be a Hilbert space, let C be a nonempty closed convex subset of H , and let $\{S_n\}$ and $\{T_n\}$ be sequences of nonexpansive self-mappings of C . Suppose that $F = F(\{S_n\}) \cap F(\{T_n\})$ is nonempty, both $\{S_n\}$ and $\{T_n\}$ satisfy the conditions (R) and (Z), and $\{S_n\}$ or $\{T_n\}$ is a strongly nonexpansive sequence. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $[0, 1]$ such that*

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty \quad \text{and} \quad 0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1.$$

Let $x, u \in C$ and let $\{x_n\}$ be a sequence in C defined by $x_1 = x \in C$ and

$$x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n (\alpha_n u + (1 - \alpha_n) T_n x_n)$$

for all $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to $P_F u$.

For $x_1, u, v \in C$, let $\{u_n\}$, $\{v_n\}$, $\{y_n\}$ and $\{x_n\}$ be the sequences defined by

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0, \\ y_n = \delta_n u_n + (1 - \delta_n) v_n, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n (\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (1.1)$$

where $f : C \rightarrow C$ is a contractive mapping with $\alpha \in (0, \frac{1}{2})$ and $\{S_n\}, \{T_n\}$ are sequences of nonexpansive mappings, one of them is a strongly nonexpansive sequence.

In this paper, inspired and motivated by [10] and [9], we prove that a strong convergence theorem of the iterative scheme $\{x_n\}$ defined by (1.1) converges strongly to $z = P_{\mathbb{F}}f(z)$, where $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{S_n\}) \cap F(\{T_n\})$, under the conditions (R) and (Z) and suitable conditions of $\{r_n\}, \{s_n\}, \{\alpha_n\}, \{\beta_n\}$ and $\{\delta_n\}$.

2 Preliminaries

In this section, we need the following lemmas to prove our main result in the next section.

Lemma 2.1 (See [11]) *Given $x \in H$ and $y \in C$. Then $P_C x = y$ if and only if the following inequality holds:*

$$\langle x - y, y - z \rangle \geq 0, \quad \forall z \in C.$$

Lemma 2.2 (See [12]) *Let $\{s_n\}$ be a sequence of nonnegative real numbers satisfying*

$$s_{n+1} = (1 - \alpha_n)s_n + \alpha_n\beta_n, \quad \forall n \geq 0,$$

where $\{\alpha_n\}, \{\beta_n\}$ satisfy the conditions

- (1) $\{\alpha_n\} \subset [0, 1], \sum_{n=1}^{\infty} \alpha_n = \infty$;
- (2) $\limsup_{n \rightarrow \infty} \beta_n \leq 0$ or $\sum_{n=1}^{\infty} |\alpha_n\beta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} s_n = 0$.

Lemma 2.3 (See [13]) *Let $\{x_n\}$ and $\{z_n\}$ be bounded sequences in a Banach space X , and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose that*

$$x_{n+1} = \beta_n x_n + (1 - \beta_n)z_n$$

for all integers $n \geq 0$ and

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Then $\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0$.

Lemma 2.4 (See [14]) *Let C be a closed convex subset of a strictly convex Banach space E . Let $\{T_n : n \in \mathbb{N}\}$ be a sequence of nonexpansive mappings on C . Suppose that $\bigcap_{n=1}^{\infty} F(T_n)$ is nonempty. Let $\{\lambda_n\}$ be a sequence of positive numbers with $\sum_{n=1}^{\infty} \lambda_n = 1$. Then a mapping S on C defined by*

$$S(x) = \sum_{n=1}^{\infty} \lambda_n T_n x$$

for all $x \in C$ is well defined, nonexpansive and $F(S) = \bigcap_{n=1}^{\infty} F(T_n)$ holds.

Lemma 2.5 (See [4]) *Let C be a nonempty closed convex subset of a Hilbert space H , and let $G : C \times C \rightarrow \mathbb{R}$ satisfy*

- (A1) $G(x, x) = 0, \forall x \in C$;
 (A2) G is monotone, i.e., $G(x, y) + G(y, x) \leq 0, \forall x, y \in C$;
 (A3) $\forall x, y, z \in C, \lim_{t \rightarrow 0^+} G(tz + (1-t)x, y) \leq G(x, y)$;
 (A4) $\forall x \in C, y \mapsto G(x, y)$ is convex and lower semicontinuous.

For $x \in H$ and $r > 0$, define a mapping $S_r : H \rightarrow C$ as follows:

$$S_r(x) = \left\{ z \in C : G(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}.$$

Then S_r is well defined and the following hold:

- (1) S_r is single-valued;
- (2) S_r is firmly nonexpansive, i.e., $\|S_r(x) - S_r(y)\|^2 \leq \langle S_r(x) - S_r(y), x - y \rangle, \forall x, y \in H$;
- (3) $F(S_r) = EP(G)$;
- (4) $EP(G)$ is closed and convex.

Lemma 2.6 (See [11]) (*Demiclosedness principle*) Assume that T is a nonexpansive self-mapping of a closed convex subset C of a Hilbert space H . If T has a fixed point, then $I - T$ is demiclosed. That is, whenever $\{x_n\}$ is a sequence in C weakly converging to some $x \in C$ and the sequence $\{(I - T)x_n\}$ converges strongly to some y , it follows that $(I - T)x = y$. Here, I is the identity mapping of H .

Lemma 2.7 Let H be a real Hilbert space. Then, for all $x, y \in H$,

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle.$$

Lemma 2.8 (See [10]) Let H be a Hilbert space, let C be a nonempty subset of H , and let $\{S_n\}$ and $\{T_n\}$ be the sequences of nonexpansive self-mappings of C . Suppose that $\{S_n\}$ and $\{T_n\}$ satisfy the condition (R) and that $\{T_n y : n \in \mathbb{N}, y \in D\}$ is bounded for any bounded subset D of C . Then $\{S_n T_n\}$ satisfies the condition (R).

Lemma 2.9 (See [1]) Let H be a Hilbert space, let C be a nonempty subset of H , and let $\{S_n\}$ and $\{T_n\}$ be the sequences of nonexpansive self-mappings of C . Suppose that $\{S_n\}$ or $\{T_n\}$ is a strongly nonexpansive sequence and that $\tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\})$ is nonempty. Then $\tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\}) = \tilde{F}(\{S_n T_n\})$.

3 Main result

Theorem 3.1 Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_1 and F_2 be two bifunctions from $C \times C$ into \mathbb{R} satisfying (A1)-(A4), respectively, and let $\{S_n\}$ and $\{T_n\}$ be sequences of nonexpansive self-mappings of C with $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{S_n\}) \cap F(\{T_n\}) \neq \emptyset$. Let $\{T_n\}$ or $\{S_n\}$ be a sequence of strongly nonexpansive mappings, and let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}, \{u_n\}, \{v_n\}$ be sequences generated by $x_1, u, v \in C$ and

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0, \\ y_n = \delta_n u_n + (1 - \delta_n) v_n, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (3.1)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (iii) $\sum_{n=0}^{\infty} |r_{n+1} - r_n|, \sum_{n=0}^{\infty} |s_{n+1} - s_n| < \infty$;
- (iv) $\lim_{n \rightarrow \infty} \delta_n = \delta \in (0, 1)$;
- (v) $\{S_n\}$ and $\{T_n\}$ satisfy the conditions R and Z.

Then the sequences $\{x_n\}, \{u_n\}, \{v_n\}, \{y_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$.

Proof Let $v \in \mathbb{F}$. From the definition of x_n , we have

$$\begin{aligned}
 \|x_{n+1} - v\| &= \|\beta_n(x_n - v) + (1 - \beta_n)(S_n(\alpha_n f(T_n y_n) + (1 - \alpha_n)T_n y_n) - v)\| \\
 &\leq \beta_n \|x_n - v\| + (1 - \beta_n) \|\alpha_n f(T_n y_n) + (1 - \alpha_n)T_n y_n - v\| \\
 &\leq \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\| + (1 - \alpha_n) \|T_n y_n - v\|) \\
 &\leq \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - f(v)\| + \alpha_n \|f(v) - v\| \\
 &\quad + (1 - \alpha_n) \|T_n y_n - v\|) \\
 &\leq \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \alpha \|y_n - v\| + \alpha_n \|f(v) - v\| \\
 &\quad + (1 - \alpha_n) \|y_n - v\|) \\
 &= \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \|f(v) - v\| \\
 &\quad + (1 - \alpha_n(1 - \alpha)) \|y_n - v\|).
 \end{aligned} \tag{3.2}$$

From Lemma 2.5 and (3.1), we have $EP(F_1) = F(S_{r_n})$, $EP(F_2) = F(S_{s_n})$, $S_{r_n}x_n = u_n$ and $S_{s_n}x_n = v_n$. By $v \in \mathbb{F}$ and the nonexpansiveness of S_{r_n} and S_{s_n} , we have

$$\begin{aligned}
 \|y_n - v\| &= \|\delta_n(u_n - v) + (1 - \delta_n)(v_n - v)\| \\
 &\leq \delta_n \|u_n - v\| + (1 - \delta_n) \|v_n - v\| \\
 &= \delta_n \|S_{r_n}x_n - v\| + (1 - \delta_n) \|S_{s_n}x_n - v\| \\
 &\leq \|x_n - v\|.
 \end{aligned} \tag{3.3}$$

Substituting (3.3) into (3.2), we have

$$\begin{aligned}
 \|x_{n+1} - v\| &\leq \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \|f(v) - v\| \\
 &\quad + (1 - \alpha_n(1 - \alpha)) \|y_n - v\|) \\
 &\leq \beta_n \|x_n - v\| + (1 - \beta_n) (\alpha_n \|f(v) - v\| \\
 &\quad + (1 - \alpha_n(1 - \alpha)) \|x_n - v\|) \\
 &= \beta_n \|x_n - v\| + (1 - \beta_n) \alpha_n \|f(v) - v\| \\
 &\quad + (1 - \beta_n) (1 - \alpha_n(1 - \alpha)) \|x_n - v\| \\
 &= (1 - \beta_n) \alpha_n \|f(v) - v\| + (1 - \alpha_n(1 - \beta_n)(1 - \alpha)) \|x_n - v\| \\
 &\leq \max \left\{ \|x_n - v\|, \frac{\|f(v) - v\|}{1 - \alpha} \right\}.
 \end{aligned}$$

By induction we can conclude that $\{x_n\}$ is bounded and so are $\{u_n\}$, $\{v_n\}$, $\{y_n\}$. Next, we show that $\tilde{F}(\{S_n A_n\}) = \tilde{F}(\{S_n\})$ and $\tilde{F}(\{A_n T_n\}) = \tilde{F}(\{T_n\})$, where $A_n = \alpha_n f + (1 - \alpha_n)I$.

Let $\{z_n\}$ be a bounded sequence in C . From the nonexpansiveness of S_n , we have

$$\|S_n A_n z_n - S_n z_n\| \leq \|A_n z_n - z_n\| = \alpha_n \|f(z_n) - z_n\|. \quad (3.4)$$

From (3.4) and $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} \|S_n A_n z_n - S_n z_n\| = 0. \quad (3.5)$$

Let $\{z_n\} \in \tilde{F}(\{S_n A_n\})$, then we have

$$\|z_n - S_n z_n\| \leq \|z_n - S_n A_n z_n\| + \|S_n A_n z_n - S_n z_n\|.$$

From (3.5), we have

$$\lim_{n \rightarrow \infty} \|z_n - S_n z_n\| = 0,$$

which implies that $\{z_n\} \in \tilde{F}(\{S_n\})$. It follows that

$$\tilde{F}(\{S_n A_n\}) \subseteq \tilde{F}(\{S_n\}). \quad (3.6)$$

Let $\{z_n\} \in \tilde{F}(\{S_n\})$, then we have

$$\|z_n - S_n A_n z_n\| \leq \|z_n - S_n z_n\| + \|S_n z_n - S_n A_n z_n\|.$$

From (3.5), we have

$$\lim_{n \rightarrow \infty} \|z_n - S_n A_n z_n\| = 0,$$

which implies that $\{z_n\} \in \tilde{F}(\{S_n A_n\})$. It follows that

$$\tilde{F}(\{S_n\}) \subseteq \tilde{F}(\{S_n A_n\}). \quad (3.7)$$

From (3.6) and (3.7), we have

$$\tilde{F}(\{S_n\}) = \tilde{F}(\{S_n A_n\}). \quad (3.8)$$

Let $\{z_n\}$ be a bounded sequence in C , then we have $\{T_n z_n\}$ is bounded and so is $\{f(T_n z_n)\}$. Since

$$\|A_n T_n z_n - T_n z_n\| = \alpha_n \|f(T_n z_n) - T_n z_n\|$$

and $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} \|A_n T_n z_n - T_n z_n\| = 0. \quad (3.9)$$

Let $\{z_n\} \in \widetilde{F}(\{A_n T_n\})$, then we have

$$\|z_n - T_n z_n\| \leq \|z_n - A_n T_n z_n\| + \|A_n T_n z_n - T_n z_n\|.$$

From (3.9), we have

$$\lim_{n \rightarrow \infty} \|z_n - T_n z_n\| = 0,$$

which implies that

$$\{z_n\} \in \widetilde{F}(\{T_n\}).$$

It follows that

$$\widetilde{F}(\{A_n T_n\}) \subseteq \widetilde{F}(\{T_n\}). \quad (3.10)$$

Let $\{z_n\} \in \widetilde{F}(\{T_n\})$, then we have

$$\|z_n - A_n T_n z_n\| \leq \|z_n - T_n z_n\| + \|T_n z_n - A_n T_n z_n\|.$$

From (3.9), we have

$$\lim_{n \rightarrow \infty} \|z_n - A_n T_n z_n\| = 0,$$

which implies that

$$\{z_n\} \in \widetilde{F}(\{A_n T_n\}).$$

It follows that

$$\widetilde{F}(\{T_n\}) \subseteq \widetilde{F}(\{A_n T_n\}). \quad (3.11)$$

From (3.10) and (3.11), we have

$$\widetilde{F}(\{T_n\}) = \widetilde{F}(\{A_n T_n\}). \quad (3.12)$$

Next, we show that

$$\widetilde{F}(\{S_n A_n T_n\}) = \widetilde{F}(\{S_n\}) \cap \widetilde{F}(\{T_n\}).$$

Since \mathbb{F} is nonempty, from (3.8), (3.12), we have

$$\widetilde{F}(\{S_n A_n\}) \cap \widetilde{F}(\{T_n\}) = \widetilde{F}(\{S_n\}) \cap \widetilde{F}(\{T_n\}) \neq \emptyset \quad (3.13)$$

and

$$\widetilde{F}(\{S_n\}) \cap \widetilde{F}(\{A_n T_n\}) = \widetilde{F}(\{S_n\}) \cap \widetilde{F}(\{T_n\}) \neq \emptyset. \quad (3.14)$$

Suppose that $\{S_n\}$ is a strongly nonexpansive sequence. From (3.14) and Lemma 2.9, we have

$$\tilde{F}(\{S_n A_n T_n\}) = \tilde{F}(\{S_n\}) \cap \tilde{F}(\{A_n T_n\}) = \tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\}). \quad (3.15)$$

On the other hand, suppose that $\{T_n\}$ is a strongly nonexpansive sequence. From (3.13) and Lemma 2.9, we have

$$\tilde{F}(\{S_n A_n T_n\}) = \tilde{F}(\{S_n A_n\}) \cap \tilde{F}(\{T_n\}) = \tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\}). \quad (3.16)$$

From (3.16) and (3.15), we have $\tilde{F}(\{S_n A_n T_n\}) = \tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\})$. Next, we show that $\{A_n\}$ and $\{S_n A_n T_n\}$ satisfy the condition (R). It is easy to see that A_n is a nonexpansive mapping for every $n \in \mathbb{N}$ and that $\{A_n y : n \in \mathbb{N}, y \in D\}$ is bounded, where D is a bounded subset of C . Let $y \in D$, then we have

$$\begin{aligned} \|A_{n+1}y - A_n y\| &= \|\alpha_{n+1}f(y) + (1 - \alpha_{n+1})y - \alpha_n f(y) - (1 - \alpha_n)y\| \\ &\leq |\alpha_{n+1} - \alpha_n| \|f(y)\| + |\alpha_{n+1} - \alpha_n| \|y\|. \end{aligned}$$

From the condition (i), we have

$$\lim_{n \rightarrow \infty} \sup_{y \in D} \|A_{n+1}y - A_n y\| = 0.$$

It follows that $\{A_n\}$ satisfies the condition (R). From Lemma 2.8, we have that $\{S_n A_n\}$ satisfies the condition (R). From the nonexpansiveness of T_n and $\mathbb{F} \neq \emptyset$, we have $\{T_n y : n \in \mathbb{N}, y \in D\}$ is bounded for any bounded subset D of C . From Lemma 2.8, we have that $\{S_n A_n T_n\}$ satisfies the condition (R).

Next, we show that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (3.17)$$

Put

$$x_{n+1} = \beta_n x_n + (1 - \beta_n) w_n, \quad (3.18)$$

where $w_n = S_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n)$. From the definition of w_n , we have

$$\begin{aligned} \|w_{n+1} - w_n\| &= \|S_{n+1} A_{n+1} T_{n+1} y_{n+1} - S_n A_n T_n y_n\| \\ &\leq \|S_{n+1} A_{n+1} T_{n+1} y_{n+1} - S_n A_n T_n y_{n+1}\| + \|S_n A_n T_n y_{n+1} - S_n A_n T_n y_n\| \\ &\leq \sup_{y \in D} \|S_{n+1} A_{n+1} T_{n+1} y - S_n A_n T_n y\| + \|y_{n+1} - y_n\|, \end{aligned} \quad (3.19)$$

where D is a bounded subset of C . Besides, we have

$$\begin{aligned} \|y_{n+1} - y_n\| &= \|\delta_{n+1} u_{n+1} + (1 - \delta_{n+1}) v_{n+1} - \delta_n u_n - (1 - \delta_n) v_n\| \\ &= \|\delta_{n+1} u_{n+1} - \delta_{n+1} u_n + \delta_{n+1} u_n - (1 - \delta_{n+1}) v_n + (1 - \delta_{n+1}) v_n\| \end{aligned}$$

$$\begin{aligned}
 & + (1 - \delta_{n+1})v_{n+1} - \delta_n u_n - (1 - \delta_n)v_n \| \\
 & = \| \delta_{n+1}(u_{n+1} - u_n) + (\delta_{n+1} - \delta_n)u_n + (1 - \delta_{n+1})(v_{n+1} - v_n) + (\delta_n - \delta_{n+1})v_n \| \\
 & \leq \delta_{n+1} \|u_{n+1} - u_n\| + |\delta_{n+1} - \delta_n| \|u_n\| + (1 - \delta_{n+1}) \|v_{n+1} - v_n\| \\
 & \quad + |\delta_n - \delta_{n+1}| \|v_n\|.
 \end{aligned} \tag{3.20}$$

From (3.1) and Lemma 2.5, we have $u_n = S_{r_n} x_n$. This implies that

$$F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0 \quad \text{for all } u \in C \tag{3.21}$$

and

$$F_1(u_{n+1}, u) + \frac{1}{r_{n+1}} \langle u - u_{n+1}, u_{n+1} - x_{n+1} \rangle \geq 0 \quad \text{for all } u \in C. \tag{3.22}$$

Putting $u = u_{n+1}$ in (3.21) and $u = u_n$ in (3.22), we have

$$F_1(u_n, u_{n+1}) + \frac{1}{r_n} \langle u_{n+1} - u_n, u_n - x_n \rangle \geq 0 \tag{3.23}$$

and

$$F_1(u_{n+1}, u_n) + \frac{1}{r_{n+1}} \langle u_n - u_{n+1}, u_{n+1} - x_{n+1} \rangle \geq 0. \tag{3.24}$$

Summing up the last two inequalities and using (A2), we obtain

$$\left\langle u_{n+1} - u_n, \frac{u_n - x_n}{r_n} - \frac{u_{n+1} - x_{n+1}}{r_{n+1}} \right\rangle \geq 0.$$

This implies that

$$\left\langle u_{n+1} - u_n, u_n - u_{n+1} + u_{n+1} - x_n - \frac{r_n}{r_{n+1}}(u_{n+1} - x_{n+1}) \right\rangle \geq 0.$$

Hence,

$$\begin{aligned}
 \|u_{n+1} - u_n\|^2 & \leq \left\langle u_{n+1} - u_n, u_{n+1} - x_n - \frac{r_n}{r_{n+1}}(u_{n+1} - x_{n+1}) \right\rangle \\
 & = \left\langle u_{n+1} - u_n, u_{n+1} - x_{n+1} + x_{n+1} - x_n - \frac{r_n}{r_{n+1}}(u_{n+1} - x_{n+1}) \right\rangle \\
 & = \left\langle u_{n+1} - u_n, x_{n+1} - x_n + \left(1 - \frac{r_n}{r_{n+1}}\right)(u_{n+1} - x_{n+1}) \right\rangle \\
 & \leq \|u_{n+1} - u_n\| \left(\|x_{n+1} - x_n\| + \frac{1}{r_{n+1}} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\| \right) \\
 & \leq \|u_{n+1} - u_n\| \left(\|x_{n+1} - x_n\| + \frac{1}{a} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\| \right).
 \end{aligned}$$

Then we have

$$\|u_{n+1} - u_n\| \leq \|x_{n+1} - x_n\| + \frac{1}{a} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\|. \tag{3.25}$$

From (3.1) and Lemma 2.5, we have $v_n = S_{s_n}x_n$. This implies that

$$F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0 \quad \text{for all } v \in C.$$

By using the same method as (3.25), we have

$$\|v_{n+1} - v_n\| \leq \|x_{n+1} - x_n\| + \frac{1}{a} |s_{n+1} - s_n| \|v_{n+1} - x_{n+1}\|. \quad (3.26)$$

Substituting (3.25) and (3.26) into (3.20), we have

$$\begin{aligned} \|y_{n+1} - y_n\| &\leq \delta_{n+1} \|u_{n+1} - u_n\| + |\delta_{n+1} - \delta_n| \|u_n\| + (1 - \delta_{n+1}) \|v_{n+1} - v_n\| \\ &\quad + |\delta_n - \delta_{n+1}| \|v_n\| \\ &\leq \delta_{n+1} \left(\|x_{n+1} - x_n\| + \frac{1}{a} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\| \right) \\ &\quad + (1 - \delta_{n+1}) \left(\|x_{n+1} - x_n\| + \frac{1}{a} |s_{n+1} - s_n| \|v_{n+1} - x_{n+1}\| \right) \\ &\quad + 2M |\delta_n - \delta_{n+1}| \\ &\leq \|x_{n+1} - x_n\| + \frac{1}{a} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\| \\ &\quad + \frac{1}{a} |s_{n+1} - s_n| \|v_{n+1} - x_{n+1}\| + 2M |\delta_n - \delta_{n+1}|, \end{aligned} \quad (3.27)$$

where $M = \sup_{n \in \mathbb{N}} \{\|u_n\|, \|v_n\|\}$. Substituting (3.27) into (3.19), we have

$$\begin{aligned} \|w_{n+1} - w_n\| &\leq \sup_{y \in D} \|S_{n+1}A_{n+1}T_{n+1}y - S_nA_nT_ny\| + \|y_{n+1} - y_n\| \\ &\leq \sup_{y \in D} \|S_{n+1}A_{n+1}T_{n+1}y - S_nA_nT_ny\| + \|x_{n+1} - x_n\| \\ &\quad + \frac{1}{a} |r_{n+1} - r_n| \|u_{n+1} - x_{n+1}\| \\ &\quad + \frac{1}{a} |s_{n+1} - s_n| \|v_{n+1} - x_{n+1}\| + 2M |\delta_n - \delta_{n+1}|. \end{aligned} \quad (3.28)$$

From (3.28), the conditions (iii), (iv) and $\{S_nA_nT_n\}$ satisfying the condition (R), we have

$$\limsup_{n \rightarrow \infty} (\|w_{n+1} - w_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (3.29)$$

From Lemma 2.3 and the definition of x_n , we have

$$\lim_{n \rightarrow \infty} \|x_n - w_n\| = 0. \quad (3.30)$$

From the definition of x_n , we have

$$x_{n+1} - x_n = (1 - \beta_n)(w_n - x_n). \quad (3.31)$$

From (3.30), (3.31) and the condition (ii), we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0.$$

Next, we show that

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0.$$

From the definition of y_n , we have

$$\|y_n - x_n\| \leq \delta_n \|u_n - x_n\| + (1 - \delta_n) \|v_n - x_n\|. \quad (3.32)$$

Next, we show that

$$\lim_{n \rightarrow \infty} \|u_n - x_n\| = \lim_{n \rightarrow \infty} \|v_n - x_n\| = 0.$$

Let $v \in \mathbb{F}$. From the definition of x_n , we have

$$\begin{aligned} \|x_{n+1} - v\|^2 &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) \|S_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n) - v\|^2 \\ &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) \|\alpha_n (f(T_n y_n) - v) + (1 - \alpha_n)(T_n y_n - v)\|^2 \\ &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\|^2 + (1 - \alpha_n) \|T_n y_n - v\|^2) \\ &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\|^2 + (1 - \alpha_n) \|y_n - v\|^2) \\ &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\|^2 \\ &\quad + (1 - \alpha_n) (\delta_n \|u_n - v\|^2 + (1 - \delta_n) \|v_n - v\|^2)). \end{aligned} \quad (3.33)$$

From the firm nonexpansiveness of S_{r_n} and $u_n = S_{r_n} x_n$, we have

$$\begin{aligned} \|u_n - v\|^2 &= \|S_{r_n} x_n - S_{r_n} v\|^2 \\ &\leq \langle u_n - v, x_n - v \rangle \\ &= \frac{1}{2} (\|u_n - v\|^2 + \|x_n - v\|^2 - \|u_n - x_n\|^2). \end{aligned}$$

It implies that

$$\|u_n - v\|^2 \leq \|x_n - v\|^2 - \|u_n - x_n\|^2. \quad (3.34)$$

Since S_{s_n} is a firmly nonexpansive mapping and $v_n = S_{s_n} x_n$, by using the same method as (3.34), we have

$$\|v_n - v\|^2 \leq \|x_n - v\|^2 - \|v_n - x_n\|^2. \quad (3.35)$$

Substituting (3.34), (3.35) into (3.33), we have

$$\begin{aligned} \|x_{n+1} - v\|^2 &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\|^2 \\ &\quad + (1 - \alpha_n) (\delta_n \|u_n - v\|^2 + (1 - \delta_n) \|v_n - v\|^2)) \\ &\leq \beta_n \|x_n - v\|^2 + (1 - \beta_n) (\alpha_n \|f(T_n y_n) - v\|^2 \\ &\quad + (1 - \alpha_n) (\delta_n (\|x_n - v\|^2 - \|u_n - x_n\|^2) \end{aligned}$$

$$\begin{aligned}
& + (1 - \delta_n)(\|x_n - v\|^2 - \|v_n - x_n\|^2)) \\
& = \beta_n \|x_n - v\|^2 + (1 - \beta_n)(\alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad + (1 - \alpha_n)(\delta_n \|x_n - v\|^2 - \delta_n \|u_n - x_n\|^2 \\
& \quad + (1 - \delta_n)\|x_n - v\|^2 - (1 - \delta_n)\|v_n - x_n\|^2)) \\
& = \beta_n \|x_n - v\|^2 + (1 - \beta_n)(\alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad + (1 - \alpha_n)(\|x_n - v\|^2 - \delta_n \|u_n - x_n\|^2 \\
& \quad - (1 - \delta_n)\|v_n - x_n\|^2)) \\
& = \beta_n \|x_n - v\|^2 + (1 - \beta_n)\alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad + (1 - \alpha_n)(1 - \beta_n)(\|x_n - v\|^2 - \delta_n \|u_n - x_n\|^2 \\
& \quad - (1 - \delta_n)\|v_n - x_n\|^2) \\
& = \beta_n \|x_n - v\|^2 + (1 - \beta_n)\alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad + (1 - \alpha_n)(1 - \beta_n)\|x_n - v\|^2 - \delta_n(1 - \alpha_n)(1 - \beta_n)\|u_n - x_n\|^2 \\
& \quad - (1 - \delta_n)(1 - \alpha_n)(1 - \beta_n)\|v_n - x_n\|^2 \\
& \leq \|x_n - v\|^2 + \alpha_n \|f(T_n y_n) - v\|^2 - \delta_n(1 - \alpha_n)(1 - \beta_n)\|u_n - x_n\|^2 \\
& \quad - (1 - \delta_n)(1 - \alpha_n)(1 - \beta_n)\|v_n - x_n\|^2.
\end{aligned} \tag{3.36}$$

From (3.36), we have

$$\begin{aligned}
\delta_n(1 - \alpha_n)(1 - \beta_n)\|u_n - x_n\|^2 & \leq \|x_n - v\|^2 - \|x_{n+1} - v\|^2 + \alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad - (1 - \delta_n)(1 - \alpha_n)(1 - \beta_n)\|v_n - x_n\|^2 \\
& \leq (\|x_n - v\| + \|x_{n+1} - v\|)\|x_{n+1} - x_n\| + \alpha_n \|f(T_n y_n) - v\|^2 \\
& \quad - (1 - \delta_n)(1 - \alpha_n)(1 - \beta_n)\|v_n - x_n\|^2 \\
& \leq (\|x_n - v\| + \|x_{n+1} - v\|)\|x_{n+1} - x_n\| \\
& \quad + \alpha_n \|f(T_n y_n) - v\|^2.
\end{aligned}$$

From the conditions (i), (ii), (iv) and (3.17), we have

$$\lim_{n \rightarrow \infty} \|u_n - x_n\| = 0. \tag{3.37}$$

By using the method as (3.37), we have

$$\lim_{n \rightarrow \infty} \|v_n - x_n\| = 0. \tag{3.38}$$

From (3.32), (3.37) and (3.38), we have

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \tag{3.39}$$

Next, we show that

$$\{y_n\} \in \tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\}). \tag{3.40}$$

Since

$$\begin{aligned}\|S_n A_n T_n y_n - y_n\| &\leq \|S_n A_n T_n y_n - x_n\| + \|x_n - y_n\| \\ &= \|w_n - x_n\| + \|x_n - y_n\|,\end{aligned}$$

from (3.30) and (3.39), we have

$$\lim_{n \rightarrow \infty} \|S_n A_n T_n y_n - y_n\| = 0.$$

Since $\{y_n\}$ is bounded, we have

$$\{y_n\} \in \tilde{F}(\{S_n A_n T_n\}). \quad (3.41)$$

Since $\tilde{F}(\{S_n A_n T_n\}) = \tilde{F}(\{S_n\}) \cap \tilde{F}(\{T_n\})$ and (3.41), we have (3.40).

Next, we show that

$$\lim_{n \rightarrow \infty} \|S_n m_n - m_n\| = 0,$$

where $m_n = \alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n$. From the definition of m_n , we have

$$\begin{aligned}\|S_n m_n - m_n\| &\leq \|S_n m_n - x_n\| + \|m_n - x_n\| \\ &= \|S_n m_n - x_n\| + \|\alpha_n (f(T_n y_n) - x_n) + (1 - \alpha_n)(T_n y_n - x_n)\| \\ &\leq \|w_n - x_n\| + \alpha_n \|f(T_n y_n) - x_n\| + (1 - \alpha_n) \|T_n y_n - x_n\| \\ &\leq \|w_n - x_n\| + \alpha_n \|f(T_n y_n) - x_n\| \\ &\quad + \|T_n y_n - y_n\| + \|y_n - x_n\|.\end{aligned}$$

From (3.39), (3.40), (3.30) and the condition (i), we have

$$\lim_{n \rightarrow \infty} \|S_n m_n - m_n\| = 0.$$

Next, we show that

$$\limsup_{n \rightarrow \infty} \langle f(z) - z, m_n - z \rangle \leq 0,$$

where $z = P_{\mathbb{F}} f(z)$. Since $\{y_n\}$ is bounded, there exists a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ converging weakly to v , that is, $y_{n_i} \rightharpoonup v$ as $i \rightarrow \infty$. From (3.40), $\{S_n\}$ and $\{T_n\}$ satisfying the condition (Z), we have $v \in F(\{S_n\}) \cap F(\{T_n\})$.

Define the mapping $Q: C \rightarrow C$ by

$$Q(x) = \delta S_{r_n} x + (1 - \delta) S_{s_n} x \quad \text{for all } x \in C,$$

where $\lim_{n \rightarrow \infty} \delta_n = \delta \in (0, 1)$. From the nonexpansiveness of S_{r_n} , S_{s_n} and Lemma 2.4, we have

$$F(Q) = F(S_{r_n}) \cap F(S_{s_n}) = EP(F_1) \cap EP(F_2).$$

From the definitions of y_n and Q , we have

$$\begin{aligned}\|x_n - Qx_n\| &\leq \|x_n - y_n\| + \|y_n - Qx_n\| \\ &\leq \|x_n - y_n\| + \|\delta_n u_n + (1 - \delta_n)v_n - \delta S_{r_n}x_n - (1 - \delta)S_{s_n}x_n\| \\ &\leq \|x_n - y_n\| + |\delta_n - \delta|\|u_n\| + |\delta_n - \delta|\|v_n\|.\end{aligned}\quad (3.42)$$

From (3.39), (3.42) and the condition (iv), we have

$$\lim_{n \rightarrow \infty} \|x_n - Qx_n\| = 0. \quad (3.43)$$

From (3.39) and $y_{n_i} \rightarrow v$ as $i \rightarrow \infty$, we have $x_{n_i} \rightarrow v$ as $i \rightarrow \infty$. By (3.43), $x_{n_i} \rightarrow v$ as $i \rightarrow \infty$ and Lemma 2.6, we have

$$v \in F(Q) = EP(F_1) \cap EP(F_2).$$

Hence,

$$v \in EP(F_1) \cap EP(F_2) \cap F(\{S_n\}) \cap F(\{T_n\}) = \mathbb{F}. \quad (3.44)$$

By (3.40), (3.44) and the condition (i), we have

$$\begin{aligned}\limsup_{n \rightarrow \infty} \langle f(z) - z, m_n - z \rangle &= \limsup_{n \rightarrow \infty} \langle \alpha_n \langle f(z) - z, f(T_n y_n) - T_n y_n \rangle \\ &\quad + \langle f(z) - z, T_n y_n - z \rangle \rangle \\ &= \lim_{i \rightarrow \infty} \langle \alpha_{n_i} \langle f(z) - z, f(T_{n_i} y_{n_i}) - T_{n_i} y_{n_i} \rangle \\ &\quad + \langle f(z) - z, T_{n_i} y_{n_i} - z \rangle \rangle \\ &= \lim_{i \rightarrow \infty} \langle \alpha_{n_i} \langle f(z) - z, f(T_{n_i} y_{n_i}) - T_{n_i} y_{n_i} \rangle \\ &\quad + \langle f(z) - z, T_{n_i} y_{n_i} - y_{n_i} \rangle + \langle f(z) - z, y_{n_i} - z \rangle \rangle \\ &= \langle f(z) - z, v - z \rangle \leq 0.\end{aligned}$$

Finally, we show that the sequence $\{x_n\}$ converges strongly to $z = P_{\mathbb{F}}f(z)$. From the definition of $\{x_n\}$, we have

$$\begin{aligned}\|x_{n+1} - z\|^2 &= \|\beta_n(x_n - z) + (1 - \beta_n)(S_n m_n - z)\|^2 \\ &\leq \beta_n \|x_n - z\|^2 + (1 - \beta_n) \|S_n m_n - z\|^2 \\ &\leq \beta_n \|x_n - z\|^2 + (1 - \beta_n) \|m_n - z\|^2.\end{aligned}\quad (3.45)$$

Since $m_n = \alpha_n f(T_n y_n) + (1 - \alpha_n)T_n y_n$, we have

$$\begin{aligned}\|m_n - z\|^2 &= \|\alpha_n(f(T_n y_n) - z) + (1 - \alpha_n)(T_n y_n - z)\|^2 \\ &\leq (1 - \alpha_n)^2 \|T_n y_n - z\|^2 + 2\alpha_n \langle f(T_n y_n) - z, m_n - z \rangle \\ &\leq (1 - \alpha_n) \|x_n - z\|^2 + 2\alpha_n \langle f(T_n y_n) - f(z), m_n - z \rangle\end{aligned}$$

$$\begin{aligned}
 & + 2\alpha_n \langle f(z) - z, m_n - z \rangle \\
 \leq & (1 - \alpha_n) \|x_n - z\|^2 + 2\alpha_n \alpha \|x_n - z\| \|m_n - z\| \\
 & + 2\alpha_n \langle f(z) - z, m_n - z \rangle \\
 \leq & (1 - \alpha_n) \|x_n - z\|^2 + \alpha_n \alpha (\|x_n - z\|^2 + \|m_n - z\|^2) \\
 & + 2\alpha_n \langle f(z) - z, m_n - z \rangle \\
 = & (1 - \alpha_n) \|x_n - z\|^2 + \alpha_n \alpha \|x_n - z\|^2 + \alpha_n \alpha \|m_n - z\|^2 \\
 & + 2\alpha_n \langle f(z) - z, m_n - z \rangle \\
 = & (1 - \alpha_n(1 - \alpha)) \|x_n - z\|^2 + \alpha_n \alpha \|m_n - z\|^2 \\
 & + 2\alpha_n \langle f(z) - z, m_n - z \rangle.
 \end{aligned}$$

This implies that

$$\begin{aligned}
 \|m_n - z\|^2 & \leq \frac{1 - \alpha_n(1 - \alpha)}{1 - \alpha_n \alpha} \|x_n - z\|^2 + \frac{2\alpha_n}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle \\
 & = \frac{1 - \alpha_n \alpha + \alpha_n \alpha - \alpha_n(1 - \alpha)}{1 - \alpha_n \alpha} \|x_n - z\|^2 + \frac{2\alpha_n}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle \\
 & = \left(1 - \frac{\alpha_n(1 - 2\alpha)}{1 - \alpha_n \alpha}\right) \|x_n - z\|^2 + \frac{2\alpha_n}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle. \tag{3.46}
 \end{aligned}$$

Substituting (3.46) into (3.45), we have

$$\begin{aligned}
 \|x_{n+1} - z\|^2 & \leq \beta_n \|x_n - z\|^2 + (1 - \beta_n) \|m_n - z\|^2 \\
 & \leq \beta_n \|x_n - z\|^2 + (1 - \beta_n) \left(\left(1 - \frac{\alpha_n(1 - 2\alpha)}{1 - \alpha_n \alpha}\right) \|x_n - z\|^2 \right. \\
 & \quad \left. + \frac{2\alpha_n}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle \right) \\
 & \leq \beta_n \|x_n - z\|^2 + (1 - \beta_n) \left(1 - \frac{\alpha_n(1 - 2\alpha)}{1 - \alpha_n \alpha}\right) \|x_n - z\|^2 \\
 & \quad + \frac{2\alpha_n(1 - \beta_n)}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle \\
 & = \beta_n \|x_n - z\|^2 + \left((1 - \beta_n) - \frac{\alpha_n(1 - 2\alpha)(1 - \beta_n)}{1 - \alpha_n \alpha} \right) \|x_n - z\|^2 \\
 & \quad + \frac{2\alpha_n(1 - \beta_n)}{1 - \alpha_n \alpha} \langle f(z) - z, m_n - z \rangle \\
 & = \left(1 - \frac{\alpha_n(1 - 2\alpha)(1 - \beta_n)}{1 - \alpha_n \alpha}\right) \|x_n - z\|^2 \\
 & \quad + \frac{\alpha_n(1 - \beta_n)(1 - 2\alpha)}{1 - \alpha_n \alpha} \frac{2 \langle f(z) - z, m_n - z \rangle}{(1 - 2\alpha)}. \tag{3.47}
 \end{aligned}$$

Applying (3.47), the conditions (i), (ii) and Lemma 2.2, we have $\{x_n\}$ converges strongly to $z = P_{\mathbb{F}}f(z)$. From (3.39), (3.37) and (3.38), it is easy to see that $\{y_n\}$, $\{u_n\}$, $\{v_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$. This completes the proof. \square

4 Applications

In this section, we give three examples for a strongly nonexpansive sequence and prove a strong convergence theorem associated to the variational inequality problem.

Before we give three examples, we need the following definition and lemmas.

Definition 4.1 Let C be a nonempty closed convex subset of a real Hilbert space H . A mapping $A : C \rightarrow H$ is called an α -inverse strongly monotone mapping if there exists an $\alpha > 0$ such that

$$\langle x - y, Ax - Ay \rangle \geq \alpha \|Ax - Ay\|^2$$

for all $x, y \in C$.

A mapping $A : C \rightarrow H$ is called α -strongly monotone if there exists $\alpha > 0$ such that

$$\langle x - y, Ax - Ay \rangle \geq \alpha \|x - y\|^2$$

for all $x, y \in C$.

A mapping $T : C \rightarrow C$ is called a κ -strictly pseudo-contractive mapping if there is $\kappa \in [0, 1)$ such that

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + \kappa \|(I - T)x - (I - T)y\|^2 \quad (4.1)$$

for all $x, y \in C$.

Then (4.1) is equivalent to

$$\langle x - y, (I - T)x - (I - T)y \rangle \geq \frac{1 - \kappa}{2} \|(I - T)x - (I - T)y\|^2$$

for all $x, y \in C$.

The set of solutions of the variational inequality problem of the mapping $A : C \rightarrow H$ is denoted by $VI(C, A)$, that is,

$$VI(C, A) = \{x \in C : \langle y - x, Ax \rangle \geq 0, \forall y \in C\}.$$

Let $A, B : C \rightarrow H$ be two mappings. In 2013, Kangtunyakarn [15] modified $VI(C, A)$ as follows:

$$VI(C, aA + (1 - a)B) = \{x \in C : \langle y - x, (aA + (1 - a)B)x \rangle \geq 0, \forall y \in C \text{ and } a \in (0, 1)\}.$$

Remark 4.1 If $T : C \rightarrow C$ is a κ -strictly pseudo-contractive mapping with $F(T) \neq \emptyset$, then $(I - T)$ is a $\frac{1 - \kappa}{2}$ -inverse strongly monotone mapping and $F(T) = VI(C, I - T)$.

Lemma 4.2 (See [16]) Let H be a Hilbert space, let C be a nonempty closed convex subset of H , and let A be a mapping of C into H . Let $u \in C$. Then, for $\lambda > 0$,

$$u = P_C(I - \lambda A)u \iff u \in VI(C, A),$$

where P_C is the metric projection of H onto C .

Lemma 4.3 (See [15]) *Let C be a nonempty closed convex subset of a real Hilbert space H , and let $A, B : C \rightarrow H$ be α and β -inverse strongly monotone mappings, respectively, with $\alpha, \beta > 0$ and $VI(C, A) \cap VI(C, B) \neq \emptyset$. Then*

$$VI(C, aA + (1-a)B) = VI(C, A) \cap VI(C, B), \quad \forall a \in (0, 1). \quad (4.2)$$

Furthermore, if $0 < \gamma < 2\eta$, where $\eta = \min\{\alpha, \beta\}$, we have $I - \gamma(aA + (1-a)B)$ is a nonexpansive mapping.

Example 4.4 Let $T : C \rightarrow C$ be a κ -strictly pseudo-contractive mapping with $F(T) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < 1 - \kappa \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n(I - T))$. Then $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z).

Proof Since T is a κ -strictly pseudo-contractive mapping, then $I - T$ is $\frac{1-\kappa}{2}$ -inverse strongly monotone. From Example 4.3 in [10], we have $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). \square

Example 4.5 Let $A, B : C \rightarrow H$ be α, β -inverse strongly monotone mappings, respectively, with $\bar{\gamma} = \min\{\alpha, \beta\}$ and $VI(C, A) \cap VI(C, B) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < 2\bar{\gamma} \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n D)$, where $D = aA + (1-a)B$ for all $a \in (0, 1)$. Then $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z).

Proof Let $x, y \in C$, then we have

$$\begin{aligned} \langle x - y, Dx - Dy \rangle &= \langle x - y, (aA + (1-a)B)x - (aA + (1-a)B)y \rangle \\ &\geq a\langle x - y, Ax - Ay \rangle + (1-a)\langle x - y, Bx - By \rangle \\ &\geq a\alpha\|Ax - Ay\|^2 + (1-a)\beta\|Bx - By\|^2 \\ &\geq \bar{\gamma}(\|aAx + (1-a)Bx - aAy - (1-a)By\|^2) \\ &\geq \bar{\gamma}\|Dx - Dy\|^2. \end{aligned}$$

Then D is a $\bar{\gamma}$ -inverse strongly monotone mapping. From Example 4.3 in [10], we have that $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). \square

Example 4.6 Let $A : C \rightarrow H$ be an α -strongly monotone and L -Lipschitzian mapping with $VI(C, A) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < \frac{2\alpha}{L^2} \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n A)$. Then $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z).

Proof Let $x, y \in C$, then we have

$$\begin{aligned}\langle x - y, Ax - Ay \rangle &\geq \alpha \|x - y\|^2 \\ &\geq \frac{\alpha}{L^2} \|Ax - Ay\|^2.\end{aligned}$$

Then A is an $\frac{\alpha}{L^2}$ -inverse strongly monotone mapping. From Example 4.3 in [10], we have that $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). \square

Example 4.7 (See [10]) Let $\{R_n\}$ be a sequence of nonexpansive mappings of C into itself having a common fixed point, and let $\{\mu_n\}$ be a sequence in $[0, 1]$. For each $n \in \mathbb{N}$, a W -mapping [17] T_n generated by R_n, R_{n-1}, \dots, R_1 and $\mu_n, \mu_{n-1}, \dots, \mu_1$ is defined as follows:

$$\begin{aligned}U_{n,n} &= \mu_n R_n + (1 - \mu_n)I, \\ U_{n,n-1} &= \mu_{n-1} R_{n-1} U_{n,n} + (1 - \mu_{n-1})I, \\ U_{n,n-2} &= \mu_{n-2} R_{n-2} U_{n,n-1} + (1 - \mu_{n-2})I, \\ &\vdots \\ U_{n,k} &= \mu_k R_k U_{n,k+1} + (1 - \mu_k)I, \\ &\vdots \\ U_{n,2} &= \mu_2 R_2 U_{n,3} + (1 - \mu_2)I, \\ T_n &= U_{n,1} = \mu_1 R_1 U_{n,2} + (1 - \mu_1)I.\end{aligned}$$

If $0 < \mu_1 \leq 1$ and $0 < \mu_n \leq b$, for all $n \geq 2$ and $0 < b < 1$, then $\{T_n\}$ satisfies the conditions (R) and (Z).

By using our main result and these three examples, we obtain the following results.

Theorem 4.8 Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_1 and F_2 be two bifunctions from $C \times C$ into \mathbb{R} satisfying (A1)-(A4), respectively. Let $T : C \rightarrow C$ be a κ -strictly pseudo-contractive mapping with $F(T) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < 1 - \kappa \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n(I - T))$. Let $\{R_n\}$ be a sequence of nonexpansive mappings of C into itself having a common fixed point, and let $\{\mu_n\}$ be a sequence in $[0, 1]$. For each $n \in \mathbb{N}$, W_n is a W -mapping generated by R_n, R_{n-1}, \dots, R_1 and $\mu_n, \mu_{n-1}, \dots, \mu_1$. Assume that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{R_n\}) \cap F(T) \neq \emptyset$. Let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}$, $\{u_n\}$, $\{v_n\}$ be sequences generated by

$x_1, u, v \in C$ and

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0, \\ y_n = \delta_n u_n + (1 - \delta_n) v_n, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) W_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (4.3)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (iii) $\sum_{n=0}^{\infty} |r_{n+1} - r_n|, \sum_{n=0}^{\infty} |s_{n+1} - s_n| < \infty$;
- (iv) $\lim_{n \rightarrow \infty} \delta_n = \delta \in (0, 1)$.

Then the sequences $\{x_n\}, \{u_n\}, \{v_n\}, \{y_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$.

Proof From Example 4.4, we have $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). From Lemma 4.2, we have $F(T_n) = F(P_C(I - \lambda_n(I - T))) = VI(C, I - T) = F(T)$ for all $n \in \mathbb{N}$. It implies that $F(\{T_n\}) = F(T)$. From [18], we have $F(\{W_n\}) = F(\{R_n\})$. It follows that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{W_n\}) \cap F(\{T_n\}) \neq \emptyset$. From Example 4.7, we have $\{W_n\}$ is a nonexpansive sequence satisfying the conditions (R) and (Z). By Theorem 3.1, we can conclude the desired result. \square

Theorem 4.9 Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_1 and F_2 be two bifunctions from $C \times C$ into \mathbb{R} satisfying (A1)-(A4), respectively. Let $A, B : C \rightarrow H$ be α, β -inverse strongly monotone mappings, respectively, with $\bar{\gamma} = \min\{\alpha, \beta\}$ and $VI(C, A) \cap VI(C, B) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < 2\bar{\gamma} \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n D)$, where $D = aA + (1 - a)B$ for all $a \in (0, 1)$. Let $\{R_n\}$ be a sequence of nonexpansive mappings of C into itself having a common fixed point, and let $\{\mu_n\}$ be a sequence in $[0, 1]$. For each $n \in \mathbb{N}$, W_n is a W -mapping generated by R_n, R_{n-1}, \dots, R_1 and $\mu_n, \mu_{n-1}, \dots, \mu_1$. Assume that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{R_n\}) \cap VI(C, A) \cap VI(C, B) \neq \emptyset$. Let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}, \{u_n\}, \{v_n\}$ be sequences generated by $x_1, u, v \in C$ and

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0, \\ y_n = \delta_n u_n + (1 - \delta_n) v_n, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) W_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (4.4)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;

- (iii) $\sum_{n=0}^{\infty} |r_{n+1} - r_n|, \sum_{n=0}^{\infty} |s_{n+1} - s_n| < \infty$;
- (iv) $\lim_{n \rightarrow \infty} \delta_n = \delta \in (0, 1)$.

Then the sequences $\{x_n\}, \{u_n\}, \{v_n\}, \{y_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$.

Proof From Example 4.5, we have $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). From Lemmas 4.2 and 4.3, we have $F(T_n) = F(P_C(I - \lambda_n D)) = VI(C, D) = VI(C, A) \cap VI(C, B)$ for all $n \in \mathbb{N}$. It implies that $F(\{T_n\}) = VI(C, A) \cap VI(C, B)$. From [18], we have $F(\{W_n\}) = F(\{R_n\})$. It follows that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{W_n\}) \cap F(\{T_n\}) \neq \emptyset$. From Example 4.7, we have $\{W_n\}$ is a nonexpansive sequence satisfying the conditions (R) and (Z). By Theorem 3.1, we can conclude the desired result. \square

Theorem 4.10 *Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_1 and F_2 be two bifunctions from $C \times C$ into \mathbb{R} satisfying (A1)-(A4), respectively. Let $A : C \rightarrow H$ be an α -strongly monotone and L -Lipschitzian mapping with $VI(C, A) \neq \emptyset$. Let $\{\lambda_n\}$ be a sequence of positive real numbers such that*

$$0 < \inf_{n \in \mathbb{N}} \lambda_n \leq \sup_{n \in \mathbb{N}} \lambda_n < \frac{2\alpha}{L^2} \quad \text{and} \quad \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) = 0,$$

and let $\{T_n\}$ be a sequence of mappings defined by $T_n = P_C(I - \lambda_n A)$. Let $\{R_n\}$ be a sequence of nonexpansive mappings of C into itself having a common fixed point, and let $\{\mu_n\}$ be a sequence in $[0, 1]$. For each $n \in \mathbb{N}$, W_n is a W -mapping generated by R_n, R_{n-1}, \dots, R_1 and $\mu_n, \mu_{n-1}, \dots, \mu_1$. Assume that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{R_n\}) \cap VI(C, A) \neq \emptyset$. Let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}, \{u_n\}, \{v_n\}$ be sequences generated by $x_1, u, v \in C$ and

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ F_2(v_n, v) + \frac{1}{s_n} \langle v - v_n, v_n - x_n \rangle \geq 0, \\ y_n = \delta_n u_n + (1 - \delta_n) v_n, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) W_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (4.5)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (iii) $\sum_{n=0}^{\infty} |r_{n+1} - r_n|, \sum_{n=0}^{\infty} |s_{n+1} - s_n| < \infty$;
- (iv) $\lim_{n \rightarrow \infty} \delta_n = \delta \in (0, 1)$.

Then the sequences $\{x_n\}, \{u_n\}, \{v_n\}, \{y_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$.

Proof From Example 4.6, we have $\{T_n\}$ is a strongly nonexpansive sequence satisfying the conditions (R) and (Z). From Lemma 4.2, we have $F(T_n) = F(P_C(I - \lambda_n A)) = VI(C, A)$ for all $n \in \mathbb{N}$. It implies that $F(\{T_n\}) = VI(C, A)$. From [18], we have $F(\{W_n\}) = F(\{R_n\})$. It follows that $\mathbb{F} = EP(F_1) \cap EP(F_2) \cap F(\{W_n\}) \cap F(\{T_n\}) \neq \emptyset$. From Example 4.7, we have $\{W_n\}$ is a nonexpansive sequence satisfying the conditions (R) and (Z). By Theorem 3.1, we can conclude the desired result. \square

Theorem 4.11 *Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_1 be a bifunction from $C \times C$ into \mathbb{R} satisfying (A1)-(A4), and let $\{S_n\}$ and $\{T_n\}$ be sequences of nonexpansive self-mappings of C with $\mathbb{F} = EP(F_1) \cap F(\{S_n\}) \cap F(\{T_n\}) \neq \emptyset$. Let $\{T_n\}$ or $\{S_n\}$ be a sequence of strongly nonexpansive mappings, and let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}, \{u_n\}$ be sequences generated by $x_1, u \in C$ and*

$$\begin{cases} F_1(u_n, u) + \frac{1}{r_n} \langle u - u_n, u_n - x_n \rangle \geq 0, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n(\alpha_n f(T_n u_n) + (1 - \alpha_n) T_n u_n), \quad \forall n \geq 1, \end{cases} \quad (4.6)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (iii) $\sum_{n=0}^{\infty} |r_{n+1} - r_n| < \infty$;
- (iv) $\{S_n\}$ and $\{T_n\}$ satisfy the conditions R and Z .

Then the sequences $\{x_n\}, \{u_n\}$ converge strongly to $z = P_{\mathbb{F}}f(z)$.

Proof Put $F_1 \equiv F_2$, $s_n = r_n$ and $u_n = v_n$. From Theorem 3.1, we can conclude the desired conclusion. \square

The following result can be obtained from Theorem 3.1. We, therefore, omit the proof.

Theorem 4.12 *Let H be a Hilbert space, let C be a nonempty closed convex subset of H . Let F_i be bifunctions from $C \times C$ into \mathbb{R} , for every $i = 1, 2, \dots, N$, satisfying (A1)-(A4), and let $\{S_n\}$ and $\{T_n\}$ be sequences of nonexpansive self-mappings of C with $\mathbb{F} = \bigcap_{i=1}^N EP(F_i) \cap F(\{S_n\}) \cap F(\{T_n\}) \neq \emptyset$. Let $\{T_n\}$ or $\{S_n\}$ be a sequence of strongly nonexpansive mappings, and let $f : C \rightarrow C$ be a contractive mapping with $\alpha \in (0, \frac{1}{2})$. Let $\{x_n\}, \{u_n\}, \{v_n\}$ be sequences generated by $x_1, u^i \in C$, for every $i \in 1, 2, \dots, N$, and*

$$\begin{cases} F_i(u_n^i, u^i) + \frac{1}{r_n^i} \langle u - u_n^i, u_n^i - x_n \rangle \geq 0, \\ y_n = \sum_{i=1}^N \delta_n^i u_n^i, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n(\alpha_n f(T_n y_n) + (1 - \alpha_n) T_n y_n), \quad \forall n \geq 1, \end{cases} \quad (4.7)$$

where $\{\alpha_n\}, \{\beta_n\} \in [0, 1]$, $\{r_n\}, \{s_n\} \in (a, b) \in [0, 1]$. Assume that the following conditions hold:

- (i) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (ii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (iii) $\sum_{n=0}^{\infty} |r_{n+1}^i - r_n^i| < \infty$, $\forall i = 1, 2, \dots, N$;
- (iv) $\sum_{i=1}^N \delta_n^i = 1$;
- (v) $\lim_{n \rightarrow \infty} \delta_n^i = \delta^i \in (0, 1)$, $\forall i = 1, 2, \dots, N$;
- (vi) $\{S_n\}$ and $\{T_n\}$ satisfy the conditions R and Z .

Then the sequences $\{x_n\}, \{y_n\}$ and $\{u_n^i\}$, for every $i = 1, 2, \dots, N$, converge strongly to $z = P_{\mathbb{F}}f(z)$.

Competing interests

The author declares that they have no competing interests.

Acknowledgements

This research was supported by the Research Administration Division of King Mongkut's Institute of Technology Ladkrabang.

Received: 29 August 2012 Accepted: 9 July 2013 Published: 22 July 2013

References

1. Aoyama, K, Kimura, Y, Takahashi, W, Toyoda, M: On a strongly nonexpansive sequence in Hilbert spaces. *J. Nonlinear Convex Anal.* **8**, 471-489 (2007)
2. Aoyama, K: An iterative method for fixed point problems for sequences of nonexpansive mappings. In: *Fixed Point Theory and Applications*, pp. 1-7. Yokohama Publ., Yokohama (2010)
3. Blum, E, Oettli, W: From optimization and variational inequalities to equilibrium problems. *Math. Stud.* **63**(1-4), 123-145 (1994)
4. Combettes, PL, Hirstoaga, SA: Equilibrium programming in Hilbert spaces. *J. Nonlinear Convex Anal.* **6**(1), 117-136 (2005)
5. Kangtunyakarn, A: Iterative methods for finding common solution of generalized equilibrium problems and variational inequality problems and fixed point problems of a finite family of nonexpansive mappings. *Fixed Point Theory Appl.* **2010**, Article ID 836714 (2010). doi:10.1155/2010/836714
6. Kangtunyakarn, A: Hybrid algorithm for finding common elements of the set of generalized equilibrium problems and the set of fixed point problems of strictly pseudocontractive mapping. *Fixed Point Theory Appl.* **2011**, Article ID 274820 (2011). doi:10.1155/2011/274820
7. Cholamjiak, W, Suantai, S: A hybrid method for a countable family of multivalued maps, equilibrium problems, and variational inequality problems. *Discrete Dyn. Nat. Soc.* **2010**, Article ID 349158 (2010). doi:10.1155/2010/349158
8. Takahashi, W, Zembayashi, K: Strong convergence theorem by a new hybrid method for equilibrium problems and relatively nonexpansive mappings. *Fixed Point Theory Appl.* **2008**, Article ID 528476 (2008). doi:10.1155/2008/528476
9. Takahashi, S, Takahashi, W: Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces. *J. Math. Anal. Appl.* **331**(1), 506-515 (2007)
10. Aoyama, K, Kimura, Y: Strong convergence theorems for strongly nonexpansive sequences. *Appl. Math. Comput.* **217**, 7537-7545 (2011)
11. Browder, FE: Convergence of approximants to fixed points of nonexpansive nonlinear mappings in Banach space. *Arch. Ration. Mech. Anal.* **24**, 82-89 (1967)
12. Xu, HK: An iterative approach to quadratic optimization. *J. Optim. Theory Appl.* **116**(3), 659-678 (2003)
13. Suzuki, T: Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals. *J. Math. Anal. Appl.* **305**(1), 227-239 (2005)
14. Bruck, RE: Properties of fixed point sets of nonexpansive mappings in Banach spaces. *Trans. Am. Math. Soc.* **179**, 251-262 (1973)
15. Kangtunyakarn, A: Convergence theorem of κ -strictly pseudocontractive mapping and a modification of generalized equilibrium problems. *Fixed Point Theory Appl.* **2012**, Article ID 89 (2012)
16. Takahashi, W: *Nonlinear Functional Analysis*. Yokohama Publ., Yokohama (2000)
17. Takahashi, W: Weak and strong convergence theorems for families of nonexpansive mappings and their applications. In: *Proceedings of Workshop on Fixed Point Theory Kazimierz Dolny*, pp. 277-292 (1997)
18. Atsushiba, S, Takahashi, W: Strong convergence theorems for a finite family of nonexpansive mappings and applications. *Indian J. Math.* **41**, 435-453 (1999)

doi:10.1186/1687-1812-2013-193

Cite this article as: Kangtunyakarn: Convergence theorems for a system of equilibrium problems and fixed point problems of a strongly nonexpansive sequence. *Fixed Point Theory and Applications* 2013 **2013**:193.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com